

琉球大学学術リポジトリ

Uncertainty principle for proper time and mass

メタデータ	言語: 出版者: 公開日: 2007-06-25 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/608

Uncertainty principle for proper time and mass

Shoju Kudaka

Department of Physics, University of the Ryukyus, Okinawa, Japan

Shuichi Matsumoto^{a)}

Department of Mathematics, University of the Ryukyus, Okinawa, Japan

(Received 6 July 1998; accepted for publication 19 November 1998)

We review Bohr's reasoning in the Bohr–Einstein debate on the photon box experiment. The essential point of his reasoning leads us to an uncertainty relation between the proper time and the rest mass of the clock. It is shown that this uncertainty relation can be derived if only we take the fundamental point of view that the proper time should be included as a dynamic variable in the Lagrangian describing the system of the clock. Some problems and some positive aspects of our approach are then discussed. © 1999 American Institute of Physics.
[S0022-2488(99)02103-9]

I. INTRODUCTION

In various arguments about time, perhaps the most spectacular is the Einstein–Bohr debate on the photon box experiment.^{1,2} Their concern in the debate was Heisenberg's time–energy uncertainty relation. However, Bohr's reasoning reveals, as shown in the following, an uncertainty relation between the proper time and the rest mass of a clock. In fact, his essential point was simply that the very act of weighing a clock, according to general relativity, interferes with the rate of the clock.

In order to review Bohr's reasoning, we consider an experiment in which we measure the rest mass of a clock. We assume, of course, that the clock keeps its own proper time.

Following Einstein's stratagem, we try to weigh the clock by suspending it with a spring. That is to say, if the spring stretches by the length l , we can calculate the mass m of the clock from the relation

$$kl = mg,$$

where g is the gravitational acceleration and k is a constant characterizing the spring.

Assume that a scale is fixed to the spring support, and that we read the length l on it with an accuracy Δq . Then the determination of l involves a minimum latitude Δp in the momentum of the clock, related to Δq by the equation $\Delta q \Delta p \approx h$. Let t be the time interval in which we read the length l . (We should note that t is measured by a clock other than the suspended clock.) Then we cannot determine the force exerted by the gravitational field on the clock to a finer accuracy than $\Delta p/t$. Therefore we cannot determine the mass m to a finer accuracy than Δm given by the relation

$$\frac{\Delta p}{t} \approx g \Delta m. \quad (1)$$

Now, according to general relativity theory, a clock, when displaced in the direction of the gravitational force by an amount Δq , changes its rate in such a way that its reading in the course of a time interval t differs by an amount $\Delta \tau$ given by the relation

^{a)}Electronic mail: shuichi@edu.u-ryukyu.ac.jp

$$\frac{\Delta\tau}{t} = \frac{g\Delta q}{c^2}. \quad (2)$$

By combining (1), (2) and the relation $\Delta q \Delta p \approx h$, we see, therefore, that there is an uncertainty relation,

$$c^2 \Delta m \Delta\tau \approx h \quad (3)$$

between the rest mass m and the proper time τ of the clock.

The relativistic redshift formula (2) was, of course, essential in Bohr's reasoning above. The more essential it seems to be, however, the stronger the apprehension we feel that the uncertainty relation (3) may fail if we can think of a weighing procedure not resorting to any interaction between the clock and the gravitational field. We check one such case in the following.

Assume that the clock has been brought to rest after being charged with an electric charge e , and that a uniform electric field \mathcal{E} is then switched on. After a short time t , we measure the distance the clock has moved. (Again t is the time measured by a clock other than our clock in the electric field.) Then we can know the average velocity v of the clock by dividing the distance by the value of t , and we can determine the mass m of the clock by virtue of the formula

$$e\mathcal{E} = m \frac{v}{t}.$$

Assume that the determination of the distance is made with a given accuracy Δq . Then it implies a minimum latitude Δp in the momentum of the clock, where $\Delta q \Delta p \approx h$. Hence, we cannot determine the force exerted by the electric field on the clock to a finer accuracy than $\Delta p/t$. Therefore, even when the velocity v is obtained, we cannot determine the mass m to a finer accuracy than Δm given by the relation

$$\frac{\Delta p}{t} \approx \Delta m \frac{v}{t}, \quad \text{i.e. } \Delta p \approx v \Delta m. \quad (4)$$

Now, according to special relativity theory, when a clock has a speed v , its rate τ in the course of a time interval t is given by the relation

$$\tau = t \sqrt{1 - (v/c)^2}. \quad (5)$$

On the other hand, the average velocity v has an uncertainty Δv given by the relation

$$t \Delta v \approx \Delta q.$$

Correspondingly, the clock has an uncertainty in its rate τ of the order $\Delta\tau$ given by

$$\Delta\tau = t \cdot \Delta \sqrt{1 - (v/c)^2} \approx \frac{v}{c^2} t \Delta v \approx \frac{v}{c^2} \Delta q. \quad (6)$$

By combining (4), (6), and the relation $\Delta q \Delta p \approx h$, we arrive, therefore, at the same uncertainty relation,

$$c^2 \Delta m \Delta\tau \approx h,$$

as (3) obtained by Bohr's reasoning.

Thus, the uncertainty relation (3) has been confirmed for a weighing procedure that does not rely on any gravitational interaction. Moreover, in this case, the time-shift formula (5) played an essential role in place of the relativistic redshift formula.

Each of these formulas is, of course, one of the deepest and most important results in relativistic theory. The fact that these important formulas play essential roles in deriving the uncertainty relation (3) lends some confidence as to its universality.

Our objective in this article is to show the following: The uncertainty relation (3) can be derived satisfactorily only if we describe the system of the clock by using a Lagrangian that includes the proper time as a dynamic variable.

In the next section, selecting the simplest Lagrangian that is in accord with the above approach, we examine the Hamiltonian formalism of the clock. Our conclusion is that the rest energy can be considered the momentum conjugate to the proper time. In the third section, following Dirac's procedure, we quantize the system of the clock, and we obtain the same uncertainty relation as (3). Some comments then follow on our quantization.

II. LAGRANGIAN AND HAMILTONIAN FORMALISM

A gravitational field $g_{\mu\nu}$ and an electromagnetic field A_μ are assumed to be given, and we consider our clock to be one material particle moving in those fields with electric charge e .

The Lagrangian that is generally used in such a case is the following:

$$L_0 = -mc\sqrt{-g_{\mu\nu}(x)\dot{x}^\mu\dot{x}^\nu} + eA_\mu(x)\dot{x}^\mu,$$

where x^μ ($\mu=0,1,2,3$) are the variables and the overdot denotes the differential with respect to an arbitrary parameter λ . It goes without saying that m is the rest mass of the clock and that c is the speed of light.

We, however, cannot consider the proper time τ a physical quantity if we describe the system by using the Lagrangian L_0 . On the other hand, it is clear that the proper time of a clock is a measurable physical quantity. (It is why a clock is so named.) Hence, we have to find another Lagrangian that is in accord with the system of the clock.

Our first purpose in this section is to find a Lagrangian L that satisfies the following conditions: (1) The Lagrangian L has the proper time τ as a new variable, in addition to x^μ . (2) The motion equations for the variables x^μ are invariant between L and L_0 .

As a candidate, we consider the Lagrangian defined by

$$L = M(\dot{\tau} - \sqrt{-g_{\mu\nu}(x)\dot{x}^\mu\dot{x}^\nu}/c) + eA_\mu(x)\dot{x}^\mu,$$

where the dynamic variables are τ , M , and x^μ .

The Lagrange's equations of motion are as follows:

$$\dot{M} = 0, \tag{7}$$

$$\dot{\tau} = \sqrt{-g_{\mu\nu}(x)\dot{x}^\mu\dot{x}^\nu}/c, \tag{8}$$

$$\frac{d}{d\lambda} \left[\frac{M}{c} \frac{g_{\rho\mu}\dot{x}^\mu}{\sqrt{-g_{\mu\nu}(x)\dot{x}^\mu\dot{x}^\nu}} + eA_\rho(x) \right] - \frac{M}{c} \frac{g_{\mu\nu,\rho}\dot{x}^\mu\dot{x}^\nu}{2\sqrt{-g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu}} - eA_{\mu,\rho}(x)\dot{x}^\mu = 0. \tag{9}$$

The second equation (8) means that we can identify the variable τ with the proper time of this clock. Moreover, we have $d\tau/d\lambda > 0$, and therefore it is possible to change the differential with respect to λ to one with respect to τ in the third equation (9). As a result, we find that

$$\frac{d}{d\tau} \left[\frac{M}{c^2} g_{\rho\mu}\dot{x}^\mu + eA_\rho(x) \right] - \frac{M}{2c^2} g_{\mu\nu,\rho}\dot{x}^\mu\dot{x}^\nu - eA_{\mu,\rho}(x)\dot{x}^\mu = 0,$$

where the overdot denotes the differential with respect to τ . Rewriting this equation, we get

$$\frac{M}{c^2} [\ddot{x}^\rho + \Gamma^\rho_{\mu\nu} \dot{x}^\mu \dot{x}^\nu] = e f^{\rho\mu} \dot{x}_\mu, \tag{10}$$

where $\Gamma^\rho_{\mu\nu}$ and $f_{\mu\nu}$ are defined by

$$\Gamma^\rho_{\mu\nu} = \frac{1}{2} g^{\rho\sigma} (-g_{\mu\nu,\sigma} + g_{\nu\sigma,\mu} + g_{\sigma\mu,\nu}), \quad f_{\mu\nu} = A_{\nu,\mu} - A_{\mu,\nu}.$$

On the other hand, the motion equation derived from the original Lagrangian L_0 is

$$m [\ddot{x}^\rho + \Gamma^\rho_{\mu\nu} \dot{x}^\mu \dot{x}^\nu] = e f^{\rho\mu} \dot{x}_\mu. \tag{11}$$

Equation (10) is just the same as Eq. (11) if we identify M with the constant mc^2 . Equation (7) indicates that this identification is possible.

Thus, our first purpose has been achieved. Moreover, this Lagrangian L is the simplest of those that satisfy the above two conditions.

The second purpose in this section is to investigate, by using the Lagrangian L , the consequences of our assertion that the proper time should be considered a dynamic variable.

We note that it is possible to propose an argument without imposing any limitation on the fields $g_{\mu\nu}$ and A_μ . In such an argument, however, we have to handle the coordinate time $x^0 = ct$ as a dynamic variable, and then determine certain constraint conditions for the variables. A discussion of such constraints is not essential for our purpose. We therefore assume for simplicity hereafter that the fields $g_{\mu\nu}$ and A_μ are so-called static in the following sense: (1) The functions $g_{\mu\nu}$ and A_μ depend on only x^1, x^2, x^3 . (2) For $i=1,2,3$, we have $g_{i0}(=g_{0i})=0$.

Assuming the above conditions, we get

$$L = M (\dot{\tau} - \sqrt{f(x)^2 - g_{ij}(x) \dot{x}^i \dot{x}^j} / c^2) + ceA_0(x) + eA_i(x) \dot{x}^i,$$

where f is defined by $g_{00} = -f^2$ ($f > 0$). The dynamic variables are τ, M, x^i ($i=1,2,3$), and the overdot denotes the differential with respect to t .

The momentum conjugate to those variables are given by

$$p_\tau \equiv \frac{\partial L}{\partial \dot{\tau}} = M, \quad p_M \equiv \frac{\partial L}{\partial \dot{M}} = 0$$

and

$$p_i \equiv \frac{\partial L}{\partial \dot{x}^i} = \frac{M}{c^2} \frac{g_{ij} \dot{x}^j}{\sqrt{f^2 - g_{jk} \dot{x}^j \dot{x}^k} / c^2} + e \dot{A}_i.$$

We have

$$H_0 = p_\tau \dot{\tau} + p_M \dot{M} + p_i \dot{x}^i - L = f \sqrt{M^2 + c^2 g^{ij} (p_i - eA_i)(p_j - eA_j)} - ceA_0.$$

If M is replaced by mc^2 , then H_0 is identical with the Hamiltonian that is derived from the original Lagrangian L_0 . In our case, however, there exist two constraints:

$$\phi_1 \equiv M - p_\tau = 0, \quad \phi_2 \equiv p_M = 0.$$

Taking account of these constraints, we have to consider the total Hamiltonian:

$$H \equiv H_0 + u_1 \phi_1 + u_2 \phi_2,$$

where u_1 and u_2 are Lagrange's undetermined multipliers.

The multipliers u_1 and u_2 are determined in the following manner:³ Poisson's bracket of ϕ_1 and ϕ_2 is

$$\{\phi_1, \phi_2\} = 1,$$

and therefore we have

$$\dot{\phi}_1 = \{\phi_1, H\} \approx u_2,$$

$$\dot{\phi}_2 = \{\phi_2, H\} \approx -u_1 - \frac{fM}{\sqrt{M^2 + c^2 g^{ij}(p_i - eA_i)(p_j - eA_j)}},$$

where the symbol " \approx " denotes the weak equality defined by the constraints $\phi_1 = \phi_2 = 0$. Hence, the consistency conditions

$$\dot{\phi}_1 \approx 0 \quad \text{and} \quad \dot{\phi}_2 \approx 0$$

require the multipliers u_1 and u_2 to be

$$u_1 = -\frac{fM}{\sqrt{M^2 + c^2 g^{ij}(p_i - eA_i)(p_j - eA_j)}} \quad \text{and} \quad u_2 = 0,$$

which give

$$H = H_0 - \frac{fM(M - p_\tau)}{\sqrt{M^2 + c^2 g^{ij}(p_i - eA_i)(p_j - eA_j)}}. \tag{12}$$

Hamilton's canonical equations of motion are as follows:

$$\dot{\tau} = \frac{\partial H}{\partial p_\tau} = \frac{fM}{\sqrt{M^2 + c^2 g^{ij}(p_i - eA_i)(p_j - eA_j)}},$$

$$\dot{p}_\tau = -\frac{\partial H}{\partial \tau} = 0,$$

$$\dot{M} = \frac{\partial H}{\partial p_M} = 0,$$

$$\dot{p}_M = -\frac{\partial H}{\partial M} \approx 0,$$

$$\dot{x}^i = \frac{\partial H}{\partial p_i} \approx \frac{\partial H_0}{\partial p_i},$$

$$\dot{p}_i = -\frac{\partial H}{\partial x^i} \approx -\frac{\partial H_0}{\partial x^i}.$$

Defining a matrix W_{ij} by

$$W_{ij} \equiv \{\phi_i, \phi_j\} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

we can write Dirac's bracket:

$$\{A, B\}_D = \{A, B\} - \sum_{i,j=1}^2 \{A, \phi_i\} W_{ij}^{-1} \{\phi_j, B\} = \{A, B\} + \{A, \phi_1\} \{\phi_2, B\} - \{A, \phi_2\} \{\phi_1, B\}.$$

We can easily calculate Dirac's brackets between the canonical variables:

$$\{\tau, p_\tau\}_D = \{\tau, M\}_D = 1, \quad \{x^i, p_j\}_D = \delta^i_j, \quad \text{the others} = 0.$$

We are now in a position to be able to state our conclusions in this section. It is easily shown that

$$\phi_1, \phi_2, T \equiv \tau - p_M, \quad E \equiv p_\tau, \quad x^i, p_i \quad (i = 1, 2, 3)$$

are canonical variables, and, therefore, the variables T, E, x^i, p_i ($i = 1, 2, 3$) can be interpreted as canonical variables on the submanifold defined by the constraints $\phi_1 = \phi_2 = 0$. We can show also that

$$\{A, B\}_D = \frac{\partial A}{\partial T} \frac{\partial B}{\partial E} - \frac{\partial A}{\partial E} \frac{\partial B}{\partial T} + \sum_{i=1}^3 \left(\frac{\partial A}{\partial x^i} \frac{\partial B}{\partial p_i} - \frac{\partial A}{\partial p_i} \frac{\partial B}{\partial x^i} \right),$$

on the submanifold.

Since we have that

$$T = \tau \quad \text{and} \quad E = M (= mc^2)$$

on the submanifold defined by $\phi_1 = \phi_2 = 0$, it follows from the above that the rest energy mc^2 is considered the momentum conjugate to the proper time τ .

III. QUANTIZATION AND DISCUSSIONS

Thus, we have arrived at the following conclusion: If we accept the view that we should describe a clock by using a Lagrangian, which includes the proper time as a dynamic variable like the positions x^i , then we find that the rest energy $E = mc^2$ turns out to be the general momentum conjugate to the proper time, and that τ, E, x^i , and p_i are canonical variables of the system.

Since τ, E, x^i, p_i are the canonical variables, if we quantize the system by Dirac's procedure, there are corresponding operators:

$$\hat{\tau}, \hat{E}, x^i, \hat{p}_i \quad (i = 1, 2, 3),$$

which satisfy the commutation relations

$$[\hat{\tau}, \hat{E}] = [x^i, \hat{p}_i] = i\hbar. \tag{13}$$

The relation $[\hat{\tau}, \hat{E}] = i\hbar$ in (13) leads us to the uncertainty relation,

$$c^2 \Delta m \Delta \tau \geq \hbar/2, \tag{14}$$

which was argued in the Introduction to this article.

Our quantization leads to some desirable results besides the uncertainty relation (14), but at the same time gives rise to some problems.

First, we should make some comment on the problems. In our quantization, the operators $\hat{\tau}, \hat{E}, x^i$ and \hat{p}_i ($i = 1, 2, 3$) can be represented in the Hilbert space composed of square integrable functions of τ, x^1, x^2 , and x^3 . In particular, the operator \hat{E} is represented by the differential

operator $-i\hbar\partial/\partial\tau$, and, therefore, the rest energy \hat{E} cannot have any discrete spectrum. Furthermore, this Hilbert space includes some states in which the mean values of \hat{E} are negative.

The problems of the continuous mass spectrum and of the negative mass are inevitable in our formulation. The authors cannot, at present, judge whether these characteristics are desirable or not. These problems will be discussed in a subsequent paper from a rather different viewpoint.

Second, we focus our attention on some positive aspects of our quantization. We restrict ourselves, for simplicity, to the case in which the space-time metric is flat and $A_\mu=0$. Then the Hamiltonian in (12) is rather simple and the Hamiltonian operator has the form

$$\hat{H} \equiv \sqrt{\hat{E}^2 + c^2 \hat{\mathbf{p}}^2}.$$

(We omit, hereafter, the overcarets representing the operators, since there is no possibility of misunderstanding.)

For the Heisenberg representation of the operator τ ,

$$\tau(t) = e^{iHt/\hbar} \tau e^{-iHt/\hbar},$$

we find that

$$\frac{d}{d\tau} \tau(t) = \frac{i}{\hbar} e^{iHt/\hbar} [H, \tau] e^{-iHt/\hbar} = \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}}, \tag{15}$$

by virtue of

$$[\tau, H] = i\hbar \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}}.$$

Hence, we have

$$\tau(t) = \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} t + \tau. \tag{16}$$

We note that the last term of (15) is the operator that represents the time delay of the moving clock.

We can, moreover, show that

$$\frac{d}{dt} \tau(t)^2 = \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \tau(t) + \tau(t) \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} = 2 \frac{E^2}{E^2 + c^2 \mathbf{p}^2} t + \left[\frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}}, \tau \right]_+,$$

where we have used Eq. (16), and where $[A, B]_+$ denotes the anticommutator of operators A and B . Integrating this, we have

$$\tau(t)^2 = \frac{E^2}{E^2 + c^2 \mathbf{p}^2} t^2 + \left[\frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}}, \tau \right]_+ t + \tau^2.$$

Hence, the standard deviation $\Delta \tau(t)$ in a state ψ is given by

$$\begin{aligned} (\Delta \tau(t))^2 \equiv \langle \tau(t)^2 \rangle - \langle \tau(t) \rangle^2 &= \left(\left\langle \frac{E^2}{E^2 + c^2 \mathbf{p}^2} \right\rangle - \left\langle \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \right\rangle^2 \right) t^2 + \left(\left\langle \left[\frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}}, \tau \right]_+ \right\rangle \right. \\ &\quad \left. - 2 \left\langle \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \right\rangle \langle \tau \rangle \right) t + (\langle \tau^2 \rangle - \langle \tau \rangle^2), \end{aligned} \tag{17}$$

where $\langle A \rangle$ denotes the mean value of an operator A in the state ψ .

Here we must introduce some approximations: We assume that the Hamiltonian operator has a very sharp value (say \mathcal{E}) in the state ψ . This assumption seems to be natural since the clock is moving as a free particle. Under this assumption, we can approximately estimate the two terms in (17) in the following manner:

$$\begin{aligned} \left\langle \frac{E^2}{E^2 + c^2 \mathbf{p}^2} \right\rangle - \left\langle \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \right\rangle^2 &\approx \frac{1}{\mathcal{E}^2} (\langle E^2 \rangle - \langle E \rangle^2), \\ \left\langle \left[\frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \cdot \tau \right]_1 \right\rangle - 2 \left\langle \frac{E}{\sqrt{E^2 + c^2 \mathbf{p}^2}} \right\rangle \langle \tau \rangle &\approx \frac{1}{\mathcal{E}} (\langle |E, \tau|_1 \rangle - 2 \langle E \rangle \langle \tau \rangle). \end{aligned} \quad (18)$$

On the other hand, the term $\langle |E, \tau|_1 \rangle - 2 \langle E \rangle \langle \tau \rangle$ in (18) often vanishes, as it does in the case of all optimal simultaneous measurements of E and τ . (We can easily check it by setting, for example, $\tau = i\hbar \partial/\partial E$ and $\psi =$ a Gaussian function of E .) Taking this cancellation into account, we neglect the second term in (17).

Thus, we have arrived at

$$(\Delta \tau(t))^2 \approx \frac{1}{\mathcal{E}^2} (\Delta E)^2 t^2 + (\Delta \tau)^2,$$

and, by virtue of the inequality

$$\frac{1}{\mathcal{E}^2} (\Delta E)^2 t^2 + (\Delta \tau)^2 \geq \frac{2}{\mathcal{E}} \Delta \tau \Delta E t,$$

we finally have

$$(\Delta \tau(t))^2 \geq \frac{\hbar}{\mathcal{E}} t, \quad (19)$$

where we have used the uncertainty relation $\Delta \tau \Delta E \geq \hbar/2$ of (14).

When the motion of the clock is so slow that the value of \mathcal{E} is approximately equal to mc^2 , then our inequality (19) has the form

$$(\Delta \tau(t))^2 \geq \frac{\hbar}{mc^2} t, \quad (20)$$

which exactly coincides with an inequality derived by Salecker and Wigner from another point of view [see Eq. (6) in Ref. 4].

In conclusion, we should make some comment on the meaning of our results to physics.

Bohr and Rosenfeld stressed the principle that every proper theory should provide in and by itself its own means for defining the quantities with which it deals. One of the key points this principle makes is that we should analyze the means of measuring those quantities in order to argue the consistency of a physical theory. In their case, they succeeded in showing that the definition of the standard quantization of electromagnetic field is consistent in the above sense by discussing the means of measuring the classical electromagnetic field.^{5,6}

Several authors have applied this principle to the theory of relativity to find a consistent quantization of the space-time geometry. The theory deals with such quantities as the metric tensor, the curvature tensor, the covariant derivative, and connection coefficients. The measurement of the distance between two events is most fundamental in the procedures by which we measure these quantities. For this we require the concept of a clock,^{7,8} and the clock cannot be independent of the various physical laws. Thus, if the above principle should be a general feature

of physical theory, a consistent formulation of the quantization of the space–time geometry should have some inherent relation with various limitations on the accuracy of the clock resulting from the physical laws.

Various gedanken experiments on such limitations have been proposed and elaborated on for some 50 years.^{4,7–16} In many of them, however, the clock is assumed to have some structure, from which starting point the argument is developed. It seems uncertain therefore whether their results are universal or not. Moreover, different studies sometimes reach different conclusions. Our objective in the present paper was to propose an attempt to dispose of this ambiguity. We showed the following: (a) There is an uncertainty relation between the proper time and the rest mass of a clock independent of its structure [see Eq. (3)]. (b) A limitation on the accuracy of the clock is derived from the uncertainty relation in a natural way [see Eqs. (19) and (20)].

The subject raised here has been argued, despite its importance, only at the level of thought experiments. The authors are uneasy with this situation, and think that the time has come to argue it at a more positive level. We hope that the importance of this subject is recognized and that, for example, the relation (20) is verified by experiment in the near future.

¹A. Pais, 'Subtle is the Lord...' *The Science and the Life of Albert Einstein* (Oxford University Press, Oxford, 1982).

²M. Jammer, *The Philosophy of Quantum Mechanics* (Wiley, New York, 1974).

³P. A. M. Dirac, *Can. J. Math.* **2**, 129 (1950); *Proc. R. Soc. London, Ser. A* **246**, 326 (1958).

⁴H. Salecker and E. P. Wigner, *Phys. Rev.* **109**, 571 (1958).

⁵N. Bohr and L. Rosenfeld, *Mat. Fys. Medd. K. Dan. Vidensk. Selsk.* **12**, XX (1933); *Phys. Rev.* **78**, 794 (1950).

⁶L. D. Landau and R. Peierls, *Z. Phys.* **69**, 56 (1931); in *Collected Papers of Landau*, edited by D. ter Haar (Gordon and Breach, New York, 1965), pp. 40–51.

⁷E. P. Wigner, *Rev. Mod. Phys.* **29**, 255 (1957).

⁸R. F. Marzke and J. A. Wheeler, in *Gravitation and Relativity*, edited by H. Y. Chiu and W. F. Hoffman (Benjamin, New York, 1964).

⁹A. Peres and N. Rosen, *Phys. Rev.* **118**, 335 (1960).

¹⁰C. A. Mead, *Phys. Rev.* **135**, B849 (1964).

¹¹F. Károlyházy, A. Frenkel, and B. Lukács, in *Quantum Concepts in Space and Time*, edited by R. Penrose and C. J. Isham (Clarendon, Oxford, 1986).

¹²A. Charlesby, *Radiat. Phys. Chem.* **33**, 487 (1989).

¹³L. Diósi and B. Lukács, *Phys. Lett. A* **142**, 331 (1989).

¹⁴F. Károlyházy, in *Sixty-Two Years of Uncertainty*, edited by A. I. Miller (Plenum, New York, 1990).

¹⁵M. Maggiore, *Phys. Lett. B* **304**, 65 (1993).

¹⁶S. Doplicher, *Ann. Inst. Henri Poincaré Phys. Theor.* **64**, 543 (1996).