

琉球大学学術リポジトリ

Aboveground respiratory behavior of two mangrove species in Manko Wetland, Okinawa Island

メタデータ	言語: 出版者: 琉球大学21世紀COEプログラム 公開日: 2008-03-09 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/4986

PE-19 Aboveground respiratory behavior of two mangrove species in Manko Wetland, Okinawa Island

Hoque A.T.M. R.¹, R. Suwa², M. Shigeta³ and A. Hagihara²

¹ Graduate School of Engineering and Science, University of the Ryukyus

² Faculty of Science, University of the Ryukyus

³ Forest Research Institute

Plant respiration plays a vital role in global carbon balance. Aboveground nighttime respiration was measured for 56 sample plants of *Kandelia obovata* Sheue, Liu & Yong and 54 sample plants of *Bruguiera gymnorrhiza* (L.) Lamk. to investigate the size-dependence of the respiratory behavior of these mangrove plants in the field conditions. At the same time, censuses were carried out in the two different monospecific plots to investigate the frequency distribution of tree size $D_{0.1H}^2H$ (H , tree height; $D_{0.1H}$, stem diameter at 10% of H) and finally to calculate stand respiration. To measure the respiration of individual plants, the enclosed standing-tree method was adopted with a CO₂ probe (Vaisala GMP343, Finland). For both the species, the dependence of the respiration on the tree size was successfully represented by a power function of $D_{0.1H}^2H$. In case of *K. obovata*, the exponent was close to 3/4. This exponent value is supported by Enquist-group's theoretical model. On the other hand, in case of *B. gymnorrhiza* the exponent was close to 2/3. This result is supported by Ninomiya-Hozumi's findings, in which the respiration per tree is proportional to its surface area. Specific respiration rate (with respect to $D_{0.1H}^2H$) was almost constant ($t = 0.421$, $p = 0.676$) under different light conditions for *K. obovata*, whereas it significantly increased with decreasing light intensity in *B. gymnorrhiza* ($t = 3.59$, $p = 7.29 \times 10^{-4}$). Stand biomass of *K. obovata* was 113.4 Mg ha⁻¹ and the stand respiration was accounted as 1.03 Mmol CO₂ ha⁻¹ yr⁻¹ (27.7 Mg d.m. ha⁻¹ yr⁻¹). Stand respiration of *B. gymnorrhiza* was 1.67 Mmol CO₂ ha⁻¹ yr⁻¹ (45.1 Mg d.m. ha⁻¹ yr⁻¹).