

琉球大学学術リポジトリ

Moonの問題 ーランダムの変差数

メタデータ	言語: 出版者: 琉球大学教育学部 公開日: 2009-04-10 キーワード (Ja): キーワード (En): 作成者: Maehara, Hiroshi, 前原, 潤 メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/9633

A Note on Moon's Problem -- Crossings in Random Graphs

Hiroshi MAEHARA*

1. Introduction

Let G be an (abstract) simple graph. Place the vertices of G randomly on the surface of a unit sphere S so that all vertices of G are distributed independently and uniformly on S . Connect two vertices a, b by the shortest arc on S whenever $\{a, b\}$ is an edge of G . The resulting configuration is called a *random drawing of G on S* . A random drawing of G on a hemisphere H of S is defined similarly. The *crossing number* of a random drawing of G is the number of pairs of arcs that intersect each other in a point interior to both. (All 'singular' cases of special position may be ignored as they occur with probability zero.)

Moon studied the crossing number $c(K_n : S)$ in a random drawing of the complete n -graph K_n on S . In [2] he stated that the distribution of $c(K_n : S)$ is asymptotically normal as n tends to infinity. However the argument to show the asymptotic normality of $c(K_n : S)$ was incorrect [3] .

We show here that the "skewness" of the distribution of $c(K_n : S)$ tends to a positive constant as n tends to infinity. Hence the distribution of $c(K_n : S)$ is never asymptotically normal. On the other hand, it is proved that the distribution of the crossing number $c(K_n : H)$ in a random drawing of K_n on a hemisphere H is asymptotically normal. It is also shown that among all graphs G with n vertices and m edges, the expected value of the crossing number in a random drawing of G on S (or H) takes the largest value when the degrees of the vertices of G are as equal as possible.

2. Geometric probability on the sphere

We recall here some results on geometric probability on a unit sphere S for later use (see [4]). For non-antipodal points a, b of S , ab denotes the shortest arc (and its length) joining them. A subset K of S is *convex* if K is hemispherical and $ab \subset K$ for every non-antipodal a, b of K .

(2.1) The probability density function of the length $s = ab$ for two random points a, b on the unit sphere S is $(1/2) \sin s$.

(2.2) The probability that a "random great circle" intersect a convex set K of perimeter L is $L / (2\pi)$.

(2.3) The mean distance between two points on the unit hemisphere H is $4/\pi$.

(2.4) The probability that four random points on the unit hemisphere H form a convex spherical quadrilateral is $3 - 24/\pi^2$.

*Dept. of Math., Coll. of Educ., Univ. of the Ryukyus

3. A complete graph on a unit sphere

Consider a random drawing of K_n on S and let V be the vertex set of the drawing. Let $x(abcd)$ be the number of crossings in the six arcs ab, ac, ad, bc, bd, cd . Then $x(abcd)$ is a random $(0, 1)$ -variable, and the crossing number $c(K_n : S)$ is written as

$$c(K_n : S) = \Sigma x(abcd) ,$$

where the summation is taken over all 4-subsets $\{a, b, c, d\}$ of V . The conditional probability that cd crosses ab given $ab = s$ follows easily from (2.2) :

$$\text{Prob}[cd \text{ crosses } ab \mid ab = s] = s/(4\pi).$$

Then by (2.1) we have the expected values of $x(abcd)$ and $c(K_n : S)$:

$$E[x(abcd)] = 3/8, \quad E[c(K_n : S)] = \binom{n}{4}(3/8).$$

Three points a, b, c , determine three great circles of the sphere S , and they divide the surface into eight spherical triangles almost surely : the triangle T_{abc} enclosed by ab, bc, ca ; the triangles T_{ab}, T_{bc}, T_{ca} , each having one side in common with T_{abc} ; the triangles T_a, T_b, T_c , each having one point in common with T_{abc} ; and the triangle T having no point in common with T_{abc} . It is easily seen that the arcs ab and cd intersect each other if and only if the point d is in the triangle T_{ab} . Since the probability that cd crosses ab under the condition $ab = s$ is $s/(4\pi)$, we have $E[\text{area}(T_{ab}) \mid ab = s]/(4\pi) = s/(4\pi)$, where $E[\mid **]$ denotes the conditional expectation under the condition $**$. Since $x(abcd)$ takes the value 1 if and only if d falls in one of T_{ab}, T_{bc}, T_{ca} , and since $\text{area}(T_{ab}) = \text{area}(T_c), \dots, \text{area}(T_{abc}) = \text{area}(T)$, we have.

$$\begin{aligned} E[x(abcd) \mid ab = s] &= \text{Prob}[x(abcd) = 1 \mid ab = s] \\ (3.1) \quad &= E[\text{area}(T_{ab}) + \text{area}(T_{bc}) + \text{area}(T_{ca}) \mid ab = s]/(4\pi) \\ &= 1/2 - E[\text{area}(T_{abc}) \mid ab = s]/(4\pi) = 1/2 - s/(4\pi). \end{aligned}$$

Hence, for different a, b, c, d, e, f , we have

$$\begin{aligned} E[x(abcd) x(abef)] &= E[(1/2 - s/(4\pi))^2] \\ &= (5\pi^2 - 4)/(32\pi^2). \end{aligned}$$

Let $y(abcd) = x(abcd) - 3/8$. Then

$$E[y(abcd) y(cdef)] = (\pi^2 - 8)/(64\pi^2).$$

(Note that $y(abcd)$ and $y(defg)$ are mutually independent as well as $y(abcd)$ and $y(efgh)$ are.)
Hence the variance of $c(K_n)$ is

$$\begin{aligned} \sigma(n)^2 &= E[(\Sigma y(abcd))^2] = \binom{n}{4} \binom{n-4}{2} \binom{4}{2} (\pi^2 - 8)/(64\pi^2) + O(n^5) \\ &= [(\pi^2 - 8)/(2^9 \pi^2)] n^6 + O(n^5). \end{aligned}$$

4. The skewness

We want to estimate the third central moment $\mu_3(n)$ of $c(K_n : S)$ when n is large. First we consider the expected value of the product $z = x(abcd) x(defg) x(ghia)$. From (3.1) it follows that

$$E[z \mid ad=s, dg=t, ga=u] = [1/(4\pi)]^3 (2\pi-s)(2\pi-t)(2\pi-u)$$

and hence

$$E[z] = 7/2^7 - [1/(4\pi)]^3 E[(ad)(dg)(ga)].$$

Let $f_{II}(s, t, u)$ be the joint probability density function of $s=ad, t=dg, u=ga$, and let $f_{III}(s, t, u)$ be the joint probability density function of s, t, u when the random three points a, d, g are chosen independently and uniformly on a fixed hemisphere H of S . Then

$$f_{II}(s, t, u) \text{Prob}(a, d, g \in H) = f(s, t, u) \text{Prob}(\Delta adg \cap G = \phi)/2,$$

where G is the great circle bounding H , and $\text{Prob}(\Delta adg \cap G = \phi)$ is the probability that G does not cut the triangle Δadg provided that the perimeter of Δadg is $s+t+u$. Then from (2.2)

$$\text{Prob}(\Delta adg \cap G = \phi) = 1 - (s+t+u)/(2\pi).$$

Hence we have

$$f_{II}(s, t, u) = 4f(s, t, u) - [2(s+t+u)/\pi] f(s, t, u).$$

Multiplying both sides by $(s)(t)(u) = (ad)(dg)(ga)$ and integrate (in full range of s, t, u such that s, t, u form a spherical triangle), we get

$$\begin{aligned} E[(ad)(dg) \mid a, d, g \in H] &= 4E[(ad)(dg)] - (4/\pi) E[(ad)^2 (dg)] \\ &\quad - (2/\pi) E[(ad)(dg)(ga)]. \end{aligned}$$

Since $E[(ad)(dg)] = E[ad]^2 = (\pi/2)^2$ and $E[(ad)^2 dg] = E[(ad)^2] E[dg] = (\pi^3 - 4\pi)/4$, we have

$$E[(ad)(dg)(ga)] = 2\pi - (\pi/2) E[(ad)(dg) \mid a, d, d \in H].$$

On the other hand

$$E[(ad)(dg) \mid a, d, g \in H] = E[w(d)^2 \mid d \in H],$$

where $w(d) = E[(ad) \mid a \in H \text{ with } d \text{ fixed}]$. Since $w(d)$ is continuous in $d \in H$ and not constant (because: by (2.3), $E[w(d) \mid d \in H] = 4/\pi$, however, if d is the "center" of H then $w(d) = 1$ by (2.1)), we must have

$$E[w(d)^2 \mid d \in H] > E[w(d) \mid d \in H]^2 = (4/\pi)^2.$$

Thus we have

$$E[(ad)(dg)(ga)] < 2\pi - (\pi/2)(4/\pi)^2 = 2\pi - 8/\pi = 3.7367\dots$$

$$< (\pi/2)^3 = 3.8757\dots$$

and $E[z] - (3/8)^3 > 7/2^7 - [1/(4\pi)]^3 (\pi/2)^3 - (3/8)^3 = 0$. Hence

$$p := E[y(abcd) y(defg) y(ghia)] = E[z] - (3/8)^3 > 0.$$

Now it is not difficult to see that the third central moment $\mu_3(n)$ is

$$\mu_3(n) = E[\{\sum y(abcd)\}^3] = \binom{n}{9} \binom{9}{4} \binom{5}{3} (36)p + O(n^8)$$

$$= (p/8) n^9 + O(n^8).$$

Thus the skewness of $c(K_n : S)$ is

$$\mu_3(n) / \sigma(n)^3 = (p/8) [2^9 \pi^2 / (\pi^2 - 8)]^{3/2} + o(1),$$

which tends to a positive constant as n tends to infinity.

5. A complete graph on a hemisphere

Here we prove the asymptotic normality of the crossing number $c(K_n : H)$ on a hemisphere H . This is a simple application of a limit lemma proved in [1]. First we state the lemma.

Let N be the set of natural numbers and r a positive integer. Suppose that for every r -

element subset A of N , there corresponds a random variable $x(A)$ defined on a common probability space and having the same mean θ . We impose the following three conditions.

(5.1) For any finite number of r -subsets $A, B, \dots, D \subset N$, the expected value $E[x(A) \dots x(D)]$ exists, and for any bijection $\tau : N \rightarrow N$, $E[x(\tau A) \dots x(\tau D)] = E[x(A) \dots x(D)]$.

(5.2) If $(A \cup \dots \cup B) \cap (C \cup \dots \cup D) = \phi$, then

$$E[x(A) \dots x(D)] = E[x(A) \dots x(B)] E[x(C) \dots x(D)].$$

Under the condition (5.1), the covariance $cov[x(A), x(B)]$ of $x(A)$ and $x(B)$ depends only on $|A \cap B|$, the number of elements in $A \cap B$. Let $c(m) = cov[x(A), x(B)]$ if $|A \cap B| = m$. Let t be the minimum value of m such that $c(m) \neq 0$.

(5.3) If $|A \cap (B \cup \dots \cup D)| \leq t$ and $|A \cap B| < t, \dots, |A \cap D| < t$, then $E[x(A) \dots x(D)] = E[x(A)] E[x(B) \dots x(D)]$.

Note that if $t = 1$ then (5.3) automatically follows from (5.2).

LEMMA. Suppose $x(A)$ (A runs over all r -subsets of N) satisfy (5.1), (5.2), (5.3), and let $s(n)$ be the sum of $x(A)$ for all r -subsets A of $\{1, 2, \dots, n\}$. Then $[s(n) - \mu] / \sigma$ tends to the normal distribution with zero mean and unit variance as n tends to infinity, where

$$\mu = \binom{n}{r} \theta, \quad \sigma^2 = [c(t) n^{2r-t}] / \{t! [(r-t)!]^2\}.$$

Now we proceed to the proof of asymptotic normality of the distribution of $c(K_n : H)$. Consider a countably infinite number of random points on the unit hemisphere H , distributed independently and uniformly on H . Label these points by natural numbers. For any 4-subset $A = \{a, b, c, d\}$ of the natural numbers, let $x(A) = x(abcd)$, the number of crossings in six arcs ab, ac, ad, bc, bd, cd . Then $x(abcd) = 1$ if four points a, b, c, d , form a convex spherical quadrilateral, and $= 0$ otherwise. Thus $\theta := E[x(A)] = 3 - 24/\pi^2$ by (2.4), and $x(A)$'s clearly satisfy the conditions (5.1), (5.2). Furthermore, $c(K_n : H) = s(n)$, the sum of $x(A)$ for all 4-subsets of $\{1, 2, \dots, n\}$.

Let $v(a) = E[x(abcd) \mid a : \text{fixed}]$. Then as a function of a , $v(a)$ is not constant. This is seen as follows. Suppose a is fixed on the boundary of the hemisphere H , and b, c be random points on H . Then H is divided by the three great circles determined by a, b, c , into six triangles (almost surely): the triangle T_{abc} enclosed by ab, bc, ca ; the triangles T_{ab}, T_{bc}, T_{ca} , each having one side in common with T_{abc} ; and triangles T_b, T_c , each having one point in common with T_{abc} . Further, these six triangles have the same expected area, as easily seen. Since $x(abcd) = 1$ if and only if d falls in T_{ab} or T_{bc} or T_{ca} , we have $E[x(abcd)] = v(a) = 1/2 \neq \theta$. Hence, when a varies in H , $v(a)$ also varies, and hence $E[x(abcd) x(aefg)] = E[v(a)^2] > \theta^2$. Thus $c(1) = cov[x(abdd), x(aefg)] > 0$, and hence we can apply the lemma. Therefore, the distribution of $c(K_n : H)$ is asymptotically normal as $n \rightarrow \infty$.

6. Crossings in a general graph

Let G be a simple graph with n vertices and m edges. Let V be the vertex set of G . Denote by $c(G)$ the number of crossings in a random drawing of G on a unit sphere S or on a hemisphere H . We show here that the expected value of $c(G)$ is

$$(6.1) \quad E[c(G)] = (\theta/2) [m^2 + m - \sum_{a \in V} deg(a)^2] ,$$

where θ is the probability that two non-adjacent arcs ab and cd cross each other, and $deg(a)$ is the degree of the vertex a of G .

Let $\{ a, b \}$ be any edge of G . Then there are

$$m - (deg(a) + deg(b)) + 1$$

edges not incident to a or b . Hence

$$E[c(G)] = \sum [m - (deg(a) + deg(b)) + 1] \theta/2$$

(where the summation is over all edges $\{ a, b \}$ of G)

$$= (\theta/2) [m^2 + m - \sum (deg(a) + deg(b))] .$$

In the summation $\sum (deg(a) + deg(b))$, each $deg(a)$ appears exactly $deg(a)$ times. Hence

$$\sum (deg(a) + deg(b)) = \sum_{a \in V} deg(a)^2 .$$

This proves (6.1).

Let \bar{d} be the average degree of G . Then the "variance" of $deg(a)$ ($a \in V$) is $(\sum deg(a)^2)/n - (\bar{d})^2$. Therefore, from (6.1) it follows that among all graphs with n vertices and m edges, $E[c(G)]$ takes the largest value when G has the minimum variance of $deg(a)$.

References

- [1] H. Maehara, On random simplices in product distributions, J. Appl. Prob. 17, 553-558 (1980).
- [2] J. W. Moon, On the distribution of crossings in random complete graphs, SIAM J., 13, 506-510 (1965).
- [3] J. W. Moon, Erratum : On the distribution of crossings in random complete graphs, SIAM J. Appl. Math. 32 (3), 706 (1977).
- [4] L. A. Santalo, *Integral Geometry and Geometric Probability*, Addison-Wesley, Reading Massachusetts (1976).