

琉球大学学術リポジトリ

U/Ca ratio in coral skeleton as a proxy of carbonate system in coral reef

メタデータ	言語: 出版者: 琉球大学21世紀COEプログラム 公開日: 2009-04-16 キーワード (Ja): キーワード (En): 作成者: Armid, Alrum, Fahmiati, Tanri, Takaesu, Yuki, Fujimura, Hiroyuki, Higuchi, Tomihiko, Taira, Eiko, Oomori, Tamotsu メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/9872

PE-14 U/Ca ratio in coral skeleton as a proxy of carbonate system in coral reef

Alrum Armid¹, Tanri Fahmiati¹, Yuki Takaesu², Hiroyuki Fujimura², Tomihiko Higuchi¹,
Eiko Taira³, and Tamotsu Oomori²

¹Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, 903-0213,
Okinawa, Japan

²Department of Chemistry, University of the Ryukyus, Nishihara, 903-0213, Okinawa, Japan

³Aqua Culture, Nishihara, Okinawa, Japan

Carbon dioxide (CO₂) is one of the most important green house gases in the atmosphere. It is recognized that CO₂ concentration has already increased since before industrial revolution to present, owing to increase in the anthropogenic activity. In addition to the green house effect, increasing CO₂ in the atmosphere enhances the CO₂ invasion from atmosphere to ocean. Increasing the amount of CO₂ dissolved in the ocean lowers the pH, and decreases the carbonate (CO₃²⁻) ions and hence the saturation state of the major skeletal carbonate minerals. It is also possible that modifying the carbonate system of the surface ocean affects the metabolism of marine organisms (such as photosynthesis and calcification). During calcification process of marine carbonates (such as coral), some trace elements (e.g. Sr, Mg) can be incorporated into carbonate skeleton and used as a proxy of the environmental change.

The concentration of total uranium in seawater is found to be ~3 ppb and distributed uniformly along the ocean as a conservative ion. Under oxic conditions, uranium mainly occurs as the hexavalent uranyl ion, UO₂²⁺. In seawater, it forms complexes with carbonate ions such as UO₂CO₃⁰, UO₂(CO₃)₂²⁻, and UO₂(CO₃)₃⁴⁻. Because of the formation of uranyl carbonate complexes is controlled by pH or CO₃²⁻ activity, uranium content of marine carbonates is possibly used to reconstruct the carbonate ion activity change in the past.

Incorporation of uranium into coral skeleton has been investigated through incubation experiment under controlled pCO₂ at 27°C. Distribution coefficient of uranium (λ_{UO_2}) between coral skeleton and seawater varied from 0.7 to 2.7 with the change in CO₃²⁻ ion activity. The data are in agreement with those of inorganic CaCO₃ precipitation experiment ($\lambda_{\text{UO}_2} = 0.8$ to 3.7)

The mode of uranium incorporation into calcium carbonate has been considered as ionic exchange of cation; CaCO₃ + UO₂²⁺ ⇌ UO₂CO₃ + Ca²⁺ (Kitano and Oomori, 1971; Broecker and Peng, 1982) or anion; CaCO₃ + UO₂(CO₃)₂²⁻ ⇌ CaUO₂(CO₃)₂ + CO₃²⁻ (Shen and Dunbar, 1995).

Such mode of uranium incorporation and its possibility to apply to the proxy of carbonate system in coral reef is discussed.