琉球大学学術リポジトリ

A remark on theorem of Jarnik

メタデータ	言語: English 出版者: Department of Mathematical Sciences, College of Science, University of the Ryukyus 公開日: 2010-01-25 キーワード (Ja): キーワード (En): 作成者: Hinokuma, Takanori, Shiga, Hiroo, 日熊, 隆則, 志賀, 博雄 メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/15068

Ryukyu Math. J., 5(1992), 1-6

A REMARK ON THEOREM OF JARNIK

TAKANORI HINOKUMA AND HIROO SHIGA

1. Introduction

In this note we consider the Hausdorff dimension of a set arising from Diophantine approximation. Set

$$E_\alpha = \{x \in (0,1) : |x - \frac{p}{q}| < \frac{1}{q^\alpha} \text{ for infinitely many positive integers } p, q\}.$$

Then the Hausdorff dimension of E_α, denoted by $dim_H E_\alpha$, is one when $\alpha \leq 2$ by Khinchine's theorem[3]. If $\alpha > 2$, $dim_H E_\alpha = \frac{2}{\alpha}$ by Jarnik's theorem[1]. The Purpose of the present note is to show the following result:

Proposition 1: *Set*

$$S_\alpha = \{x \in (0,1) : |x - \frac{p}{q}| < \frac{|\sin q|}{q^\alpha} \textit{ for infinitely many positive integers } p, q\}.$$

Then

$$dim_H S_\alpha = \begin{cases} 1 & \textit{if } \alpha \leq 2 \\ \frac{2}{\alpha} & \textit{if } \alpha > 2. \end{cases}$$

In the case $\alpha \leq 2$, the above result is contained in Khinchine's theorem since $\int_1^\infty \frac{|\sin x|}{x^{\alpha-1}} dx = \infty$.

The outline of the proof is based on [1]. But the essential part of our estimation is changed anew to avoid discussing the distribution of the set of prime numbers p satisfying $|\sin p| > c$ for a positive constant c.

Received October 31, 1992.

2. Preliminaries

We first show the inequality $dim_H S_\alpha \leq \frac{2}{\alpha}$. For each positive integer q, we set

$$S_q = \{x \in [0,1] : |x - \frac{p}{q}| < \frac{|\sin q|}{q^\alpha}\}.$$

Then S_q consists of $q-1$ intervals of length $\frac{2|\sin q|}{q^\alpha}$ and two end intervals of length $\frac{|\sin q|}{q^\alpha}$. It is clear that $S_\alpha \subset \bigcup_{q=n}^{\infty} G_q$ for each positive integer n. Hence if we choose n so that $\frac{|\sin n|}{2n^\alpha} \leq \frac{1}{2n^\alpha} \leq \delta$, then

$$\mathcal{H}^s_\delta(S_\alpha) \leq \sum_{q=n}^{\infty}(q+1)\left(\frac{2|\sin q|}{q^\alpha}\right)^s \leq \sum_{q=n}^{\infty}(q+1)\left(\frac{2}{q^\alpha}\right)^s.$$

When $s > \frac{2}{\alpha}$, the above series is convergent. So $\mathcal{H}^s(S_\alpha) = 0$ and we have $dim_H S_\alpha \leq \frac{2}{\alpha}$.

To prove converse inequality we prepare some lemmata.

Definition 2.1: *Let $\{x_n\}$, $n = 1, 2, \cdots$ be a sequence of real numbers. The sequence $\{x_n\}$ is said to be well distrbuted mod 1 if for any pair of numbers with $0 \leq a < b \leq 1$ we can choose a number N_0 which does not depend on k such that*

$$\left|\frac{\#\{x_i \in [a,b) : i = k+1, \cdots, k+N\}}{N} - (b-a)\right| < \epsilon$$

for $N \geq N_0$, where $\#$ denotes the cardinality.

Lemma 2.2: *Let c be a constant with $0 < c < 1$. Then for any $\epsilon > 0$ there exists N_0 so that if $N \geq N_0$*

$$\left|\frac{\#\{N < q \leq 2N : |\sin q| > c\}}{N} - l_c\right| < \epsilon$$

for some positive number l_c depending only on c.

Proof: Let J_c be the subinterval of $[0, \pi]$ satisfying $|\sin x| > c$. Then we have

$$\begin{aligned}
&\#\{N < q \leq 2N : |\sin q| > c\} \\
&= \#\{N < q \leq 2N : q \in J_c \bmod \pi\} \\
&= \#\{N < q \leq 2N : \frac{q}{\pi} \in \tilde{J}_c \bmod 1\},
\end{aligned}$$

where $\tilde{J}_c$ denotes the subinterval of $[0, 1]$ such that $|\sin \pi x| > c$. Since $\{\frac{q}{\pi}\}$ $q = 1, 2, \cdots$ is well distributed by Weyl's Criterion ([2], Example 5.2), for any $\epsilon > 0$, there is a number N_0 such that if $N > N_0$

$$\left| \frac{\#\{N < q \leq 2N : \frac{q}{\pi} \in \tilde{J}_c \bmod 1\}}{n} - l_c \right| < \epsilon$$

where l_c is the length of $\tilde{J}_c$. ∎

Let $[0, 1] = E_0 \supset E_1 \supset E_2 \supset \cdots$ be a decreasing sequence of sets satisfying the following conditions:

(1) each E_k is a disjoint union of finite number of closed intervals,

(2) each interval of E_k contains at least two intervals of E_{k+1},

(3) the maximal length of intervals in E_k tends to zero as $k \to \infty$.

Then the set $F = \bigcap_{k=1}^{\infty} E_k$ is totally disconnected subset of $[0, 1]$. The following lemma is contained in [1].

Lemma 2.3: *Suppose in the above construction each interval of E_{k-1} contains at least m_k intervals of E_k $(k = 1, 2, \cdots)$ which are separated by the distance at least ϵ_k with $0 < \epsilon_{k+1} < \epsilon_k$ for each k. Then*

$$dim_H F \geq \varliminf_{k \to \infty} \frac{\log(m_1 \cdots m_{k-1})}{-\log(m_k \epsilon_k)}.$$

3. Proof of Proposition 1

Throughout this section we fix a some constant real number c with $0 < c < 1$. We translate S_α to the right by one and set

$$G_q = \{x \in (1+q^{-\alpha}, 2-q^{-\alpha}) : |x - \frac{p}{q}| < \frac{c}{q^\alpha} \text{ with reduced fraction } \frac{p}{q}\}$$

Let n be a positive integer and q_i $(i = 1, 2)$ be positive integers satisfying $n < q_1 < q_2 \leq 2n$. If $\frac{r_1}{q_1} \neq \frac{r_2}{q_2}$ then

$$\left|\frac{r_1}{q_1} - \frac{r_2}{q_2}\right| = \frac{1}{q_1 q_2}|r_1 q_2 - r_2 q_1| \geq \frac{1}{q_1 q_2} \geq \frac{1}{4n^2}$$

Hence the distance between distinct centers of any pairs of intervals in G_{q_1} and G_{q_2} is at least $\frac{1}{4n^2}$. Since these intervals have lengths at most $2cn^{-\alpha}$, the distance between any point of G_{q_1} and any point of G_{q_2} is at least

$$\frac{1}{4}n^{-2} - 2cn^{-\alpha} \geq \frac{1}{8}n^{-2}$$

provided for sufficiently large $n \geq n_0$ for some large n_0. Set

$$C_n = \{n < q \leq 2n : |\sin q| > c\}.$$

By Lemma 2.2 the cardinality of the set C_n is at least $\frac{n}{2}l_c$ if $n \geq n_0'$ for some large $n_0' \geq n_0$. For such n the set

$$H_n = \bigcup_{q \in C_n} G_q$$

consists of intervals of length at least $c(2n)^{-\alpha}$ which are separated by distance at least $\frac{1}{8}n^{-2}$. Let $I = [a, b] \subset [1, 2]$ be an interval. Applying the prime number theorem there are at least

$$\frac{1}{2}\left(\frac{bq}{\log bq} - \frac{aq}{\log aq}\right)$$

primes in the range $[aq, bq]$ for sufficiently large q. Moreover we have

$$\begin{aligned}
&\frac{1}{2}\left(\frac{bq}{\log bq} - \frac{aq}{\log aq}\right) \\
&= \frac{1}{2}\frac{(b-a)q\log q + q(b\log a - a\log b)}{(\log q + \log b)(\log q + \log a)} \\
&> \frac{1}{4}\frac{(b-a)q}{\log q} = \frac{1}{4}\frac{q|I|}{\log q}
\end{aligned}$$

for sufficiently large q. Thus there are at least

$$\frac{n|I|}{4\log n}$$

reduced fractions whose denominators are q in the range $[a, b]$ if $n < q \leq 2n$ and $n \geq n_1$ for some large $n_1 > n_0'$. Summarizing the above estimates, at least

$$\frac{n^2 l_c}{8\log n}|I|$$

intervals of H_n are contained in $I = [a, b]$ provided that $n \geq n_1$.

Let n_1 be as above and $n_k = n_{k-1}^k$ for $k = 2, 3, \cdots$. Let $E_0 = [\delta, 1]$ and for $k = 1, 2, \cdots$ let E_k consist of those intervals of H_{n_k} which are completely contained in E_{k-1}. The intervals of E_k are of lengths at least $c(2n_k)^{-\alpha}$ and are separated by distance at least $\epsilon_k = \frac{1}{8}n_k^{-2}$. From the above argument each interval of E_{k-1} contains at least m_k intervals of E_k, where

$$m_k = \frac{n_k^2 l_c}{8\log n_k}c(2n_{k-1})^{-\alpha} = \frac{K n_k^2 n_{k-1}^{-\alpha}}{\log n_k}$$

if $k \geq 2$, where $K = \frac{1}{8}2^{-\alpha}cl_c$. Note that we take $m_1 = 1$ and $m_k \geq 2$ for sufficiently large k. By Lemma 2.3

$$
\begin{aligned}
&dim_H(\bigcap_{k=1}^{\infty} E_k) \\
&\geq \varliminf_{k\to\infty} \frac{\log\left[K^{k-2}n_1^{-\alpha}(n_2\cdots n_{k-2})^{2-\alpha}n_{k-1}^2(\log n_2\cdots\log n_{k-1})^{-1}\right]}{-\log\left[Kn_{k-1}^{-\alpha}(8\log n_k)^{-1}\right]} \\
&= \varliminf_{k\to\infty} \frac{\log\left[K^{k-2}n_1^{-\alpha}(n_2\cdots n_{k-2})^{2-\alpha}(\log n_2\cdots\log n_{k-1})^{-1}\right] + 2\log n_{k-1}}{-\log(K/8) + \log k(\log n_{k-1}) + \alpha\log n_{k-1}} \\
&= \frac{2}{\alpha}
\end{aligned}
$$

since $\log n_k = n_1 k!$. If $x \in E_k \subset H_{n_k}$, then $|x - \frac{p}{q}| < \frac{c}{q^\alpha}$ and $c \leq |\sin q|$ where $n_k < q \leq 2n_k$ and $\frac{p}{q}$ is a reduced fraction. Thus $\bigcap_{k=1}^{\infty} E_k \subset S_\alpha$. Hence $dim_H S_\alpha \geq \frac{2}{\alpha}$.

References

1. K.Falconner, "Fractal Geometry," John Wiley and Sons, 1990.
2. L.Kuipers and H.Niederreiter, "Uniform Distributin of Sequences," John Wiley and Sons, 1974.
3. V.G.Sprindžuk, "Metric Theory of Diophantine Approximations," Winston Wiley, 1979.

Department of Mathematics
College of Science
University of the Ryukyus
Nishihara, Okinawa 903-01
JAPAN