# 琉球大学学術リポジトリ

# Convergence of positive linear functionals

メタデータ	言語:
	出版者: Department of Mathematics, College of
	Science, University of the Ryukyus
	公開日: 2010-03-03
	キーワード (Ja):
	キーワード (En):
	作成者: Nishishiraho, Toshihiko, 西白保, 敏彦
	メールアドレス:
	所属:
URL	http://hdl.handle.net/20.500.12000/16050

Ryukyu Math. J. 1(1988), 73-94.

#### CONVERGENCE OF POSITIVE LINEAR FUNCTIONALS

### Toshihiko NISHISHIRAHO

#### 1. Introduction

Let X be a compact Hausdorff space. Let B(X) denote the Banach lattice of all real-valued bounded functions on X, endowed with the supremum norm and the canonical order. C(X) denotes the closed sublattice of B(X) consisting of all real-valued continuous functions on X. For a given point t  $\in$  X, we denote by C<sub>t</sub>(X) the closed sublattice of B(X) consisting of all real-valued bounded functions on X which are continuous at t, and by  $\delta_t$  the point evaluation functional at t, defined by  $\delta_t(f) = f(x)$  for all  $f \in B(X)$ . Let A(X) be a linear subspace of C<sub>t</sub>(X) which contains the unit function 1<sub>x</sub> defined by 1<sub>x</sub>(y) = 1 for all  $y \in X$ .

-73-

The purpose of this paper is to establish a convergence theorem in the context of positive linear functionals on A(X). Furthermore, several applications can be provided and actually, we shall obtain a generalization of the classical Korovkin theorem (cf. [6]) for sequences of positive linear functionals on C([a, b]), where [a, b] is a bounded closed interval in the real line  $\mathbb{R}$ .

For other researches of Korovkin type convergence theorems in various directions, see, e.g., [1], [2], [3], [4], [5], [8], [10], [14], [17] and [18].

#### 2. Convergence Theorems

Let t be any fixed point of X and let  $\Phi_{\mbox{t}}$  be a non-negative function in A(X) such that

(1) 
$$\inf\{\Phi_t(x); x \in F\} > 0 \text{ for every closed subset } F$$
  
of  $X \setminus \{t\}$ .

Let  $\{L_{\alpha,\lambda}; \alpha \in D, \lambda \in \Lambda\}$  be a family of positive linear functionals on A(X), where D is a directed set and  $\Lambda$  is an index set. Let  $\xi$  be any fixed non-negative real number.

Theorem 1. If

(2) 
$$\lim_{\alpha} L_{\alpha,\lambda}(\Phi_t) = 0 \quad \text{uniformly in } \lambda \in \Lambda$$

and if there exists a function  $u \in A(X)$  such that

-74-

(3) 
$$\inf\{u(x); x \in X\} > 0$$

and

(4) 
$$\lim_{\alpha} L_{\alpha,\lambda}(u) = \xi \delta_t(u) \quad \text{uniformly in } \lambda \in \Lambda,$$

then for every  $f \in A(X)$ ,

(5) 
$$\lim_{\alpha} L_{\alpha,\lambda}(f) = \xi \delta_t(f) \quad \text{uniformly in } \lambda \in \Lambda.$$

Proof. Let  $\Psi$  be a function in A(X) with  $\Psi(t) = 0$ , and let  $\varepsilon > 0$  be given. Then there exists a neighborhood V(t) of t such that  $|\Psi(x)| < \varepsilon$  for all  $x \in V(t)$ . Let  $F = X \setminus V(t)$ , and set

 $m = \inf\{\Phi_t(x); x \in F\} \text{ and } n = \sup\{|\Psi(x)|; x \in F\}.$ In view of (1), we have m > 0, and so

$$|\Psi| < \varepsilon l_X + (n/m)\Phi_t$$

which, by the positivity and linearity of  $L_{\alpha,\lambda}$ , yields

(6) 
$$|L_{\alpha,\lambda}(\Psi)| \leq \varepsilon L_{\alpha,\lambda}(1_X) + (n/m)L_{\alpha,\lambda}(\Phi_L)$$

for all  $\alpha \in D$ ,  $\lambda \in \Lambda$ . By (3), there exists a constant C > 0 such that  $u(x) \ge C$  for every  $x \in X$ , and thus

$$L_{\alpha,\lambda}(1_X) \leq (1/C)L_{\alpha,\lambda}(u) \qquad (\alpha \in D, \lambda \in \Lambda),$$

which together with (4) shows that there exists an element  $\alpha_0 \in D$  such that

(7) 
$$\sup\{L_{\alpha,\lambda}(1_X); \alpha \geq \alpha_0, \alpha \in D, \lambda \in \Lambda\} < \infty.$$

Therefore, by (2), (6) and (7) we conclude

(8) 
$$\lim_{\alpha} L_{\alpha,\lambda}(\Psi) = 0 \quad \text{uniformly in } \lambda \in \Lambda.$$

Now we take

$$\Psi = f - (f(t)/u(t))u$$

for any f  $\epsilon$  A(X). Then  $\Psi$  belongs to A(X) and  $\Psi$ (t) = 0. Hence, (8) implies

(9) 
$$\lim_{\alpha \to \lambda} \{L_{\alpha,\lambda}(f) - \delta_t(f/u)L_{\alpha,\lambda}(u)\} = 0$$
 uniformly in  $\lambda \in \Lambda$ .

Consequently, the equality

$$L_{\alpha,\lambda}(f) - \xi \delta_{t}(f) = L_{\alpha,\lambda}(f) - \delta_{t}(f/u)L_{\alpha,\lambda}(u) + \delta_{t}(f/u)\{L_{\alpha,\lambda}(u) - \xi \delta_{t}(u)\}$$

establishes the desired result (5) by virtue of (4) and (9).

Corollary 1. If (2) is fulfilled and if  $\lim_{\alpha} L_{\alpha,\lambda}(1_X) = \xi \qquad \text{uniformly in } \lambda \in \Lambda,$ 

then (5) holds for all f  $\in$  A(X).

Corollary 2. If L is a positive linear functional on A(X) satisfying  $L(\Phi_t) = 0$ , then  $L(f) = L(1_X)\delta_t(f)$  for every  $f \in A(X)$ .

Remark 1. Theorem 1, Corollaries 1 and 2 even hold for arbitrary topological spaces X.

Let p be any fixed positive real number and let G be a subset of C(X) separating the points of X. Suppose that A(X) contains the set { $|g - g(t)1_X|^p$ ;  $g \in G$ }. For each  $g \in G$ , we define

$$\mu_{\alpha,\lambda}^{(\mathbf{p},\mathbf{t})}(g) = L_{\alpha,\lambda}(|g - g(t)\mathbf{1}_{X}|^{p}) \qquad (\alpha \in D, \lambda \in \Lambda).$$

Theorem 2. If for all  $g \in G$ ,

(10) 
$$\lim_{\alpha} \mu_{\alpha,\lambda}^{(p,t)}(g) = 0 \quad \text{uniformly in } \lambda \in \Lambda$$

and if there exists a function  $u \in A(X)$  satisfying (3) and (4), then (5) holds for every  $f \in A(X)$ .

Proof. Let h be an arbitrary function in A(X). Since the original topology on X is identical with the weak topology on X induced by G, given  $\varepsilon > 0$ , we can choose a finite subset  $\{g_1, g_2, \dots, g_m\}$  of G and a constant C > 0 such that

$$|h(x) - h(t)| < \varepsilon + C \sum_{i=1}^{m} |g_i(x) - g_i(t)|^{p}$$

for all x  $\in$  X. Thus, for any  $\alpha$   $\in$  D,  $\lambda$   $\in$   $\Lambda$  we obtain

$$\left| L_{\alpha,\lambda}(h) - h(t) L_{\alpha,\lambda}(1_{X}) \right| \leq \varepsilon L_{\alpha,\lambda}(1_{X}) + C \sum_{i=1}^{m} \mu_{\alpha,\lambda}^{(p,t)}(g_{i}),$$

which together with (7) and (10) gives

(11)  $\lim_{\alpha} \{L_{\alpha,\lambda}(h) - \delta_t(h)L_{\alpha,\lambda}(1_X)\} = 0$  uniformly in  $\lambda \in \Lambda$ .

Now, for all  $\alpha \in D$ ,  $\lambda \in \Lambda$  we have

$$L_{\alpha,\lambda}(f) - \xi \delta_{t}(f) = L_{\alpha,\lambda}(f) - \delta_{t}(f)L_{\alpha,\lambda}(1_{X})$$

+ 
$$\delta_{t}(f/u) \{ \delta_{t}(u) L_{\alpha,\lambda}(1_{X}) - L_{\alpha,\lambda}(u) \}$$

+ 
$$\delta_{t}(f/u) \{ L_{\alpha,\lambda}(u) - \xi \delta_{t}(u) \},$$

which yields the desired assertion (5) on account of (4) and (11).

Corollary 3. Suppose that A(X) contains the set  $V = \{g^i; g \in G, i = 0, 1, 2, \dots, p\}$ , where p is an arbitrary fixed even positive integer. If for all  $h \in V$ ,

$$\lim_{\alpha} L_{\alpha,\lambda}(h) = \xi \delta_t(h) \qquad \text{uniformly in } \lambda \in \Lambda,$$

then (5) holds for any f  $\in$  A(X).

Corollary 4. Let L be a positive linear functional on A(X). If  $L(|g - g(t)1_X|^p) = 0$  for each  $g \in G$ , then  $L(f) = L(1_X)\delta_t(f)$  for every  $f \in A(X)$ . Also, under the hypothesis of Corollary 3, if  $L(h) = \xi\delta_t(h)$  for all  $h \in V$ , then  $L = \xi\delta_t$  on A(X).

Let M be a linear subspace of C(X) which contains  $l_X$ . For any x  $\in$  X, we denote by  $R_x(M)$  the set of all positive linear functionals L on C(X) such that L(f) = f(x) for all f  $\in$  M and by  $\varepsilon_x$  the restriction of  $\delta_x$  to C(X). Obviously,  $\varepsilon_x$  belongs to  $R_x(M)$ . We set

$$\partial_{\mathbf{M}}(\mathbf{X}) = \{\mathbf{x} \in \mathbf{X}; \mathbf{R}_{\mathbf{x}}(\mathbf{M}) = \{\mathbf{\varepsilon}_{\mathbf{x}}\}\},\$$

which is called the Choquet boundary of X with respect to M. For any f  $\in$  B(X), we define

$$f^{\#} = \inf\{g; f \leq g, g \in M\}$$

and

$$f_{\#} = \sup\{g; g \leq f, g \in M\}$$

which are called the upper and lower M-envelopes of f, respectively. Evidently, we have

$$f_{\#} \leq f \leq f^{\#}, \quad f_{\#} = -(-f)^{\#} \quad (f \in B(X)).$$

Moreover,  $f^{\#}$  and  $f_{\#}$  are upper and lower semi-continuous functions in B(X), respectively. Also, we get

$$(f + g)^{\#} \leq f^{\#} + g^{\#}$$
 (f, g  $\in B(X)$ )

and

$$(\beta f)^{\#} = \beta f^{\#}$$
 (f  $\in B(X), \beta \in \mathbb{R}, \beta \geq 0$ ).

Note that the set

$$M^{\#}(X) = \{ f \in C(X); f^{\#} = f_{\#} \} = \{ f \in C(X); f^{\#} = f = f_{\#} \}$$

is a linear subspace of C(X) containing M. Define

$$\partial_{M}^{\#}(X) = \{ x \in X; f^{\#}(x) = f_{\#}(x) \text{ for all } f \in C(X) \}$$
$$= \{ x \in X; f^{\#}(x) = f(x) \text{ for all } f \in C(X) \}.$$

Then we have

$$\partial_{M}(X) = \partial_{M}^{\#}(X)$$

and

$$\partial_{M}(X) = X$$
 if and only if  $M^{\#}(X) = C(X)$ 

(cf. [2]).

In view of these observations and Corollaries 2 and 4, we make the following remarks.

Remark 2. (i) Let M be a linear subspace of C(X) containing  $1_X$  and  $\Phi_t$ . If  $\Phi_t(t) = 0$ , then t belongs to  $\partial_M(X)$ .

(ii) Let M be a linear subspace of C(X) which contains  $1_X$  and  $\{|g - g(t)1_X|^p; g \in G\}$ . Then t belongs to  $\partial_M(X)$ .

(iii) Let V be as in Corollary 3 and let M be a linear subspace of C(X) containing V. Then we have  $\partial_M(X) = X$ , and so  $M^{\#}(X) = C(X)$ .

Remark 3. Suppose that  $M \subset A(X)$ . If for all  $g \in M$ ,

 $\lim_{\alpha} L_{\alpha,\lambda}(g) = \xi \delta_t(g) \qquad \text{uniformly in } \lambda \in \Lambda,$ 

then for all f  $\in M^{\#}(X) \cap A(X)$ ,

$$\lim_{\alpha} L_{\alpha,\lambda}(f) = \xi \delta_{t}(f) \quad \text{uniformly in } \lambda \in \Lambda.$$

Also, the statement analogous to this result can be formulated in the context of positive linear operators of a linear subspace of C(X) into B(X). These results extend [2; Proposition 1.4] (cf. [3; Theorem 1]).

## 3. Applications

Let  $\mathbf{E}_1$  and  $\mathbf{E}_2$  be normed vector lattices with norms  $||\cdot||_{\mathbf{E}_1}$  and  $||\cdot||\mathbf{E}_2$ , respectively. Let Y be a linear

-81-

subspace of  $E_1$  and let T be a positive linear operator of Y into  $E_2$ . A subset S of Y is called a Korovkin test system (or, briefly, KTS) with respect to {T, Y,  $E_1$ ,  $E_2$ } if for any family {T<sub> $\alpha,\lambda$ </sub>;  $\alpha \in D$ ,  $\lambda \in \Lambda$ } of positive linear operators of Y into  $E_2$ , the relation

 $\lim_{\alpha} || T_{\alpha,\lambda}(g) - T(g) ||_{E_2} = 0 \quad \text{uniformly in } \lambda \in \Lambda$ 

for every g  $\in$  S implies that

$$\lim_{\alpha} || T_{\alpha,\lambda}(f) - T(f) ||_{E_2} = 0 \qquad \text{uniformly in } \lambda \in \Lambda$$

for every  $f \in Y$  (cf. [10]). In particular, if  $E_1 = E_2$  or  $E_2 = \mathbb{R}$ , then such a subset S is called a Korovkin test system with respect to T in Y.

With this notion Corollary 3 claims that V is a KTS with respect to  $\xi \delta_t$  in A(X). As an immediate consequence of Theorems 1 and 2, we have the following:

Theorem 3. Let w be a function in A(X) such that  $\inf\{w(x); x \in X\} > 0$ .

(i) Suppose that  $\Phi_t(t) = 0$ . Then  $\{w, \Phi_t\}$  is a KTS with respect to  $\xi \delta_t$  in A(X). In particular,  $\{1_X, \Phi_t\}$  is a KTS with respect to  $\xi \delta_t$  in A(X).

(ii) {w}  $\cup$  {|g - g(t)1<sub>X</sub>|<sup>p</sup>; g  $\in$  G} is a KTS with respect to  $\xi \delta_{+}$  in A(X). In particular,

 $\{1_X\} \cup \{\big|g-g(t)1_X\big|^p; \ g \in G\}$  is a KTS with respect to  $\xi \delta_t$  in A(X).

Corollary 5. Let  $\{f_1, f_2, \dots, f_m\}$  be a finite subset of A(X) and let

$$\Phi_{t}(x) = \sum_{i=1}^{m} a_{i}(t)f_{i}(x) \qquad (x \in X),$$

where each  $a_i(t)$  is a real number, such that  $\Phi_t$  is a non-negative function in B(X) satisfying (1) and  $\Phi_t(t) = 0$ . Let w be as in Theorem 3. Then {w,  $f_1$ ,  $f_2$ , ...,  $f_m$ } is a KTS with respect to  $\xi \delta_t$  in A(X). In particular, {1<sub>X</sub>,  $f_1$ ,  $f_2$ , ...,  $f_m$ } is a KTS with respect to  $\xi \delta_t$  in A(X).

We shall now mention some examples of  $\Phi_t \in C(X)$ satisfying 0 =  $\Phi_t(t) < \Phi_t(x)$  for each  $x \in X$ ,  $x \neq t$ , and so (1) is always satisfied:

(1°) Let  $\{g_1, g_2, \dots, g_r\}$  be a finite subset of C(X) such that for every  $x \in X$ ,  $x \neq t$ , there exists an integer  $j \in \{1, 2, \dots, r\}$  for which  $g_j(x) \neq g_j(t)$ . Assume that A(X) contains the set

$$G_1 = \{1_X, g_1, g_2, \dots, g_r, g_1^2, g_2^2, \dots, g_r^2\}.$$

Then with the help of the function  $\Phi_t$  given by

-83-

$$\Phi_{t} = \sum_{i=1}^{r} (g_{i} - g_{i}(t)1_{X})^{2},$$

it follows from Corollary 5 that the set

$$G_2 = \{1_X, g_1, g_2, \dots, g_r, g_1^2 + g_2^2 \dots + g_r^2\}$$

is a KTS with respect to  $\xi \delta_t$  in A(X). Note that  $G_1$  is also a KTS with respect to  $\xi \delta_t$  in A(X). Furthermore, if  $\{g_1, g_2, \dots, g_r\}$  separates the points of X and if M is a linear subspace of C(X) containing  $G_1$ , then we have  $\partial_M(X) = \partial_M^{\#}(X) = X$  and  $M^{\#}(X) = C(X)$ .

For example, we take  $X = X_r$  a compact subset of  $\mathbb{R}^r$  and let us denote by  $e_i$  (i = 1, 2, ..., r) the i-th coordinate function on  $X_r$ . Suppose that  $A(X_r)$  contains the set

$$K_1 = \{1_X, e_1, e_2, \dots, e_r, e_1^2, e_2^2, \dots, e_r^2\}.$$

Then K<sub>1</sub> and

$$K_2 = \{1_X, e_1, e_2, \dots, e_r, e_1^2 + e_2^2 + \dots + e_r^2\}$$

are Korovkin test systems with respect to  $\xi \delta_t$  in A(X<sub>r</sub>). Next, let X =  $\mathbb{T}^r$  be the r-dimensional torus and for i = 1, 2, ..., r, we define

$$(x_1, x_2, \dots, x_r) = \cos x_i \quad ((x_1, x_2, \dots, x_r) \in X)$$

and

$$s_i(x_1, x_2, \dots, x_r) = \sin x_i \quad ((x_1, x_2, \dots, x_r) \in X).$$

Assume that  $A(\mathbb{T}^r)$  contains the set

$$M_r = \{1_X, c_1, c_2, \dots, c_r, s_1, s_2, \dots, s_r\}.$$

Then M is a KTS with respect to  $\xi \delta_t$  in A(Tr). In particular, the sets

$$\{1_X, e_1, e_1^2\}$$
 and  $\{1_X, c_1, s_1\}$ 

are Korovkin test systems with respect to  $\xi \delta_t$  in  $A(X_1)$ and in  $A(\mathbb{T}^1)$ , respectively. Concerning the usual convergence, for the special case  $\xi = 1$  this is the classical result due to Korovkin (cf. [6]). For further generalizations of the Korovkin theorem from other approaches, one may consult [1], [2], [3], [4], [5], [8], [10], [14], [17] and [18].

 $(2^{\circ})$  Let (X, d) be a compact metric space. Let

$$\Phi_{+}(\mathbf{x}) = \Phi(\mathbf{d}(\mathbf{x}, t)) \qquad (\mathbf{x} \in \mathbf{X}),$$

where  $\Phi$  is a strictly increasing continuous function on [0,  $\infty$ ) with  $\Phi(0) = 0$ . For instance, the case where  $\Phi$  is given by  $\Phi(y) = y^{q}$ , q > 0, may be useful (cf. (3°), (4°)). (3°) Let X be a compact subset of a normed linear space with norm  $||\cdot||$ . Let

$$\Phi_{t}(\mathbf{x}) = ||\mathbf{x} - t||^{q}, \quad q > 0, \qquad (\mathbf{x} \in \mathbf{X}).$$

(4°) Let (H, <•, •>) be a real pre-Hilbert space and let X be a compact subset of H. Let

$$\Phi_{t}(x) = \langle x - t, x - t \rangle$$
 (x  $\in X$ ).

For any  $x \in X$ , we define

$$h(x) = \langle x, x \rangle$$
 and  $h_{+}(x) = \langle x, t \rangle$ .

Then we have the following:

Theorem 4. If h and h<sub>t</sub> belong to A(X), then  $\{1_X, h, h_t\}$  is a KTS with respect to  $\xi \delta_t$  in A(X).

Proof. This follows from Corollary 1 (cf. Theorem 3 (i)).

For example, one takes  $H = \mathbb{R}^{r}$  with the usual inner product

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + \dots + x_r y_r$$

for x =  $(x_1, x_2, \dots, x_r)$ , y =  $(y_1, y_2, \dots, y_r) \in \mathbb{R}^r$ . Then we have

$$h = \sum_{i=1}^{r} e_i^2$$
 and  $h_t = \sum_{i=1}^{r} e_i(t)e_i$ .

If A(X) contains  $K_1$  and if for i = 1, 2, ..., r,

$$\lim_{\alpha} L_{\alpha,\lambda}(e_i) = \xi \delta_t(e_i) \qquad \text{uniformly in } \lambda \in \Lambda,$$

then

$$\lim_{\alpha} L_{\alpha,\lambda}(h_t) = \xi \delta_t(h_t) \quad \text{uniformly in } \lambda \in \Lambda.$$

Consequently,  $K_2$  becomes again a KTS with respect to  $\xi \delta_t$  in A(X).

Corollaries 3 and 4, Remark 2 (ii), (iii) and Theorem 3 (ii) can be applied in the following situation (cf. [14]):

(5°) Let X be a compact subset of a real locally convex Hausdorff vector space E with its dual space E\* and G = { $\tau |_X$ ;  $\tau \in E*$ }, where  $\tau |_X$  denotes the restriction of  $\tau$  to X.

For instance, taking  $E = \mathbb{R}^r$ ,  $K_1$  becomes again a KTS with respect to  $\xi \delta_t$  in A(X).

Let  $\mathbb{N}$  denote the set of all non-negative integers. Let  $B = \{b_{\alpha,n}^{(\lambda)}; \alpha \in D, \lambda \in \Lambda, n \in \mathbb{N}\}$  be a family of non-negative real numbers with  $\Sigma_{n=0}^{\infty} b_{\alpha,n}^{(\lambda)} = 1$  for each  $\alpha \in D, \lambda \in \Lambda$ . For examples of such families, see, e.g., [12] (cf. [9], [10]). Let  $\{L_n; n \in \mathbb{N}\}$  be a sequence of positive linear functionals on A(X). For any  $f \in A(X)$ , we define

(12) 
$$L_{\alpha,\lambda}(f) = \sum_{n=0}^{\infty} b_{\alpha,n}^{(\lambda)} L_n(f)$$
 ( $\alpha \in D, \lambda \in \Lambda$ ),

which converges in  $\mathbb{R}$ . Plainly, each  $L_{\alpha,\lambda}$  is a positive linear functional on A(X), and consequently our general results can be applicable to the family  $\{L_{\alpha,\lambda}\}$ .

For this it is convenient to make the following definition: Let  $(E_1, || \cdot ||_{E_1})$  and  $(E_2, || \cdot ||_{E_2})$  be normed vector lattices. Let Y be a linear subspace of  $E_1$  and T a positive linear operator of Y into  $E_2$ . A subset S of Y is called a B-Korovkin test system (or, briefly, B-KTS) with respect to {T, Y,  $E_1, E_2$ } if for any sequence {T<sub>n</sub>;  $n \in \mathbb{N}$ } of positive linear operators of Y into  $E_2$ , the relation

$$\lim_{\alpha} \left\| \sum_{n=0}^{\infty} b_{\alpha,n}^{(\lambda)} T_{n}(g) - T(g) \right\|_{E_{2}} = 0 \quad \text{uniformly in } \lambda \in \Lambda$$

for every g  $\, \in \,$  S implies that

$$\begin{split} \lim_{\alpha} || \sum_{n=0}^{\infty} b_{\alpha,n}^{(\lambda)} T_n(f) - T(f) ||_{E_2} &= 0 \qquad \text{uniformly in } \lambda \in \Lambda \\ \text{for every } f \in Y, \text{ where it is assumed that the above series} \\ \text{converge for each } \alpha, \ \lambda, \ g \text{ and } f. \quad \text{In particular, if} \\ E_1 &= E_2 \text{ or } E_1 = \mathbb{R}, \text{ then such a subset S is called a B-KTS} \\ \text{with respect to T in Y.} \end{split}$$

With this concept we are able to derive various B-Korovkin test systems. For instance, concerning the almost convergence (F-summability) introduced by Lorentz [7], which is the case where

 $D = \mathbb{N} \setminus \{0\}, \quad \Lambda = \mathbb{N},$ 

$$b_{\alpha,n}^{(\lambda)} = 1/\alpha \qquad (\lambda \leq n < \alpha + \lambda)$$

and

$$b_{\alpha,n}^{(\lambda)} = 0 \qquad (n < \lambda, \alpha + \lambda \leq n),$$

(12) reduces to

$$L_{\alpha,\lambda}(f) = (1/\alpha) \sum_{\substack{n=\lambda \\ n=\lambda}}^{\alpha+\lambda-1} L_n(f).$$

In this case,

$$\lim_{\alpha \to \infty} L_{\alpha,\lambda}(f) = \xi \delta_t(f) \qquad \text{uniformly in } \lambda \in \mathbb{N}$$

if and only if each Banach limit of the sequence  $\{L_n(f)\}$  is equal to  $\xi \delta_t(f)$  (cf. [7; Theorem 1]).

We give now the concrete examples: Let I be the unit  $\ensuremath{\mathsf{r}}$  to the unit r-cube, i.e.,

$$I_r = \{x = (x_1, x_2, \dots, x_r) \in \mathbb{R}^r; 0 \le x_i \le 1, i = 1, 2, \dots, r\}$$

and let  $B_t(I_r)$  denote the linear subspace of  $C_t(I_r)$ 

consisting of all real-valued bounded integrable functions on  $I_r$  which are continuous at t = (y, y, ..., y), where y is an arbitrary fixed point of  $I_1$ . Let  $\{k_n; n \in \mathbb{N}\}$  be a sequence of non-negative functions in  $C(I_1)$ . For each  $n \in \mathbb{N}$ , we define the functional  $L_n$  by

$$L_{n}(f) = \int_{I} \prod_{r i=1}^{r} k_{n}(e_{i}(x))f(x)dx$$

for all  $f \in B_t(I_r)$ . Put

$$v_n^{(i)} = \int_0^1 x^i k_n(x) dx$$
 (n  $\in \mathbb{N}$ , i = 0, 1, 2).

Then  $\{v_n^{(i)}\}$  converges to  $y^i$  for i = 0, 1, 2 if and only if  $\{L_n(f)\}$  converges to  $\delta_t(f)$  for every  $f \in B_t(I_r)$ . Also, if  $\{v_n^{(0)}\}$  converges to one and if  $\{v_n^{(i)}\}$  is almost convergent to  $y^i$  for i = 1, 2, then  $\{L_n(f)\}$  is almost convergent to  $\delta_t(f)$  for all  $f \in B_t(I_r)$ .

For example, if we take

$$k_n(x) = (n + 1) x^n$$
  $(n \in \mathbb{N}, 0 \le x \le 1)$ 

and t = (1, 1, ..., 1), then for every  $f \in B_t(I_r)$ , {L<sub>n</sub>(f)} converges to  $\delta_t(f)$  and it is also almost convergent to  $\delta_t(f)$ . Next, if we take

$$k_n(x) = 2(1 - x^2)^n / \rho_n$$
 (n  $\in \mathbb{N}, 0 \leq x \leq 1$ ),

where

$$\rho_{n} = \int_{-1}^{1} (1 - x^{2})^{n} dx = \Gamma(1/2)\Gamma(n + 1)/\Gamma(n + 3/2)$$
$$= 2^{2n+1} (n!)^{2}/(2n + 1)!,$$

and t = (0, 0, ..., 0), then for all  $f \in B_t(I_r)$ ,  $\{L_n(f)\}$ is almost convergent to  $\delta_t(f)$ .

Finally, it should be remarked that under certain appropriate conditions, Theorems 1 and 2 can be recast in a quantitative form in which we are able to estimate the rate of convergence of  $\{L_{\alpha,\lambda}(f)\}$  by using a modulus of continuity of f (cf. [10], [11], [12], [13], [15], [16]). We omit the details.

#### References

- [1] D. Amir and Z. Ziegler, Korovkin shadows and Korovkin systems in C(S)-spaces, J. Math. Anal. Appl., 62(1978), 640-675.
- [2] H. Bauer, Theorems of Korovkin type for adapted spaces, Ann. Inst. Fourier, 23(1973), 245-260.
- [3] H. Berens and G. G. Lorentz, Theorems of Korovkin type for positive linear operators on Banach lattices, Approximation Theory (Proc. Int. Sympos.

-91-

Austin 1973; ed. by G. G. Lorentz), 1 - 30, Academic Press, New York - London - San Francisco, 1973.

- [4] H. Berens and G. G. Lorentz, Geometric theory of Korovkin sets, J. Approx. Theory, 15(1975), 161-189.
- [5] K. Donner, Extension of positive operators and Korovkin theorems, Lecture Notes in Math. Vol. 904, Springer-Verlag, Berlin - Heidelberg - New York, 1982.
- [6] P. P. Korovkin, Linear operators and approximation theory, Hindustan Publ. Corp., Delhi, 1960.
- [7] G. G. Lorentz, A contribution to the theory of divergent sequences, Acta Math., 80(1948), 167-190.
- [8] G. G. Lorentz, Korovkin sets (Sets of convergence), Lecture Notes, Center for Numerical Analysis, No. 59, The University of Texas at Austin, 1972.
- [9] T. Nishishiraho, Quantitative theorems on approximation processes of positive linear operators, Multivariate Approximation Theory II (Proc. Int. Conf. Math. Res. Inst. Oberwolfach 1982; ed. by W. Schempp and K. Zeller), ISNM Vol. 61, 297 – 311, Birkhäuser Verlag, Basel – Boston – Stuttgart, 1982.

- [10] T. Nishishiraho, Convergence of positive linear approximation processes, Tôhoku Math. J., 35(1983), 441-458.
- [11] T. Nishishiraho, The rate of convergence of positive linear approximation processes, Approximation Theory IV (Proc. Int. Sympos. College Station, Texas 1983; ed. by C. K. Chui, L. L. Schumaker and J. D. Ward), 635-641, Academic Press, New York-London-San Francisco, 1983.
- T. Nishishiraho, The degree of approximation by positive linear approximation processes, Bull.
   Coll. Educ., Univ. Ryukyus, 28(1985), 7-36.
- [13] T. Nishishiraho, The degree of approximation by iterations of positive linear operators, Approximation Theory V (Proc. Int. Sympos. College Station, Texas 1986; ed. by C. K. Chui, L. L. Schumaker and J. D. Ward), 507 - 510, Academic Press, New York - London - San Francisco, 1986.
- [14] T. Nishishiraho, The convergence and saturation of iterations of positive linear operators, Math. Z., 194(1987), 397-404.

-93-

- [15] T. Nishishiraho, Quantitative estimates for approximation by positive linear operators, Bull. Coll. Sci., Univ. Ryukyus, 45(1987), 1-18.
- [16] T. Nishishiraho, The order of approximation by positive linear operators, Tôhoku Math. J., 40(1988), to appear.
- [17] Ju. A. Šaškin, Korovkin systems in spaces of continuous functions, Amer. Math. Soc. Transl.,
  (2) 54(1966), 125 - 144 (Izv. Acad. Nauk SSSR, Ser. Mat., 26(1962), 495 - 512).
- [18] D. E. Wulbert, Convergence of operators and Korovkin's theorem, J. Approx. Theory, 1(1968), 381-390.

Department of Mathematics (Received October 31, 1988) College of Science University of the Ryukyus Nishihara-Cho, Okinawa 903 - 01 JAPAN