

First record of *Acetes sibogae sibogae* Hansen, 1919 in Japan

Jun Fukuchi¹, Yukio Hanamura² and Hideyuki Imai^{1*}

¹Laboratory of Marine Biology and Coral Reef Studies, Faculty of Science,
University of the Ryukyus, Nishihara, Okinawa 903-0213, Japan

²Shinjo-cho 3806-40, Tanabe, Wakayama 646-0011, Japan

Abstract. The occurrence of the shallow-water sergestid shrimp *Acetes sibogae sibogae* Hansen, 1919 (Crustacea, Decapoda, Sergestidae) in Japan has been confirmed. This is the first record of this species and reflects a notable northward range expansion to Okinawa-jima Island, Ryukyu Archipelago, Japan.

Key words: new record, *Acetes sibogae sibogae*, Sergestidae, Japan

Introduction

The shallow-water sergestid shrimp *Acetes sibogae sibogae* Hansen, 1919 (Crustacea, Decapoda, Sergestidae) has been recorded in South and South-East Asian waters from India to the Philippines and the northern coast of Australia, above approximately 40°S (Omori, 1975; Xiao & Greenwood, 1993; Chan, 1998; Hanamura, 1999). Three subspecies (*A. sibogae australis*, *A. s. sibogae*, and *A. s. sibogalis*) have been recognised over this geographical range owing to their morphological characteristics (e.g., Omori, 1975; Pérez Farfante & Kensley, 1997; Hanamura, 1999), but *A. s. sibogalis* is considered a junior synonym of *A. s. sibogae* (see De Grave & Fransen, 2011). Further taxonomic analysis regarding this subspecific classification is needed (e.g., Ravindranath, 1980).

This species is important in the coastal crustacean community in Asia and has often been exploited in regional small-scale fisheries in Asian countries (Holthuis, 1980; Xiao & Greenwood, 1993; Chan, 1998; Hanamura, unpublished data).

Before the index occurrence, three *Acetes* spe-

cies, *A. japonicus* Kishinouye, 1905, *A. chinensis* Hansen, 1919, and *A. erythraeus* Nobili, 1905 were known to occur in Japan (Chihara & Murano, 1997). During a recent survey of *Acetes*, we found a considerable number of swarming shrimp in a mangrove estuary in Kokuba-gawa River (26°11'49.4"N, 127°41'11.1"E), Okinawa-jima Island, Ryukyu Archipelago, Japan. We collected samples every month between March 2017 and May 2017. These shrimp were identified as *Acetes sibogae sibogae*, which was hitherto not recorded in Japan (e.g., Hayashi, 1992).

This paper is the first record of this interesting species in Japan and provides some taxonomic information. The carapace length (cl in mm) is given to indicate the size of the shrimp. The materials examined herein have been deposited in the National Museum of Nature and Science, Tsukuba, Japan (NSMT).

Taxonomic account

Acetes sibogae sibogae Hansen, 1919

[New Japanese name: Jiboga akiami]

(figs. 1 & 2)

Restricted synonymies

Acetes sibogae Hansen, 1919: 38, pl. 3, fig. 4a-h:

*Corresponding author: imai@sci.u-ryukyu.ac.jp




Fig. 1. *Acetes sibogae sibogae* Hansen, 1919, male 3.3 mm cl (A, B) from the Kokuba-gawa River, Okinawa-jima Island, Japan: A, lateral view of whole body; B, ventral view of last abdominal segment and uropods. Scale bars: 5 mm in A; 250 µm in B.

Ravindranath, 1980: 266: figs. 8 & 9; Hanamura, 1999: 465, figs. 1 & 2.

Acetes sibogae sibogae: Omori, 1975, fig. 27.

Acetes sibogalis Achuthankutty & George, 1973: 139, figs. 1-20.

Acetes orientalis Achuthankutty & Ayyappan Nair, 1976: 233, figs. 1-19.

Materials examined.

NSMT-Cr 25571, 10 males (4.1- 4.8 mm cl), 10 females (5.0-6.5 mm cl), Kokuba-gawa River, Okinawa-jima Island, Japan, May 2017, coll. Jun Fukuchi, det. Yukio Hanamura.

Taxonomic notes

Morphological characteristics

There is a red spot on the ventral side of the last abdominal segment (Fig. 1-B). There are two red spots on the uropod, one of which is on the sympod of the uropod and the other of which is on the basal part of the endopod.

In males, the lower antennular flagellum consists of 13 segments (Fig. 2-A). There is a clasping spine and an accessory spine arising from the same segment. The following segment bears three spines. The segment opposing the tip of the clasping spine bears




Fig. 2. *Acetes sibogae sibogae* Hansen, 1919, male 4.3 mm cl (A, B) and female 6.5 mm cl (C) from the Kokuba-gawa River, Okinawa-jima Island, Japan: A, male lower antennular flagellum; B, petasma; C, thelycum. The thelycum was stained with bromophenol blue. Scale bars: 250 µm in A-1, A-2; 500 µm in B-1, B-2; 1 mm in C-1, C-2.

three to six spines, with the most common number being four or five. The capitulum of the petasma carries two falcate spines (Fig. 2-B). There are a few minute spines on the distal end of the capitulum.

In females, the lower antennular flagellum consists of 14 to 18 segments. The distal inner margin of the basis of the third pereopod terminates in a projection, and the coxa is produced into a strong blunt tooth on the distal inner margin (Fig. 2-C). The lateral margins of the third thoracic sternite are elevated; there is a pair of small protuberances on the anterior part.

Remarks

The morphological features above are consistent with those of *A. s. sibogae*. The number of falcate spines of the capitulum of the petasma differs from those in *A. s. sibogae* and *A. s. australis*, although most morphological characteristics are commonly shared between the two subspecies. Genetic analysis is required to determine the extent of differentiation between these subspecies.

Occurrence and environmental conditions

Despite searching for it in several neighbouring rivers, such as the Yuhi-gawa River, Taiho-gawa River, Oura-gawa River, and Haneji-gawa River, we found *Acetes sibogae sibogae* only in the Kokubagawa River in the Manko estuary (Ramsar Site no. 996) in Okinawa-jima Island, Japan.

The water condition of the Kokubagawa River is cloudy, whereas the other rivers are clear and highly transparent. *Acetes* consume phytoplankton, small zooplankton, fine sediments, and detritus (Xiao & Greenwood, 1993). Fine sediments and detritus cloud the river water, and water quality has a major influence on the biomass and species composition of phytoplankton and zooplankton, which *A. s. sibogae* may consume. Water quality may affect the distribution of this shrimp.

We collected all *Acetes* material between 4 pm to 7 pm, when the tide was rising. Shrimp were obtained near the muddy bottom using scoop nets. The water temperature at the sampling site ranged from 22°C to 25°C, and the salinity ranged from 22 to 33. We collected most samples when the salinity was about 27, but more detailed information is required to determine the best salinity for *A. s. sibogae*.

In mangrove estuaries on the northwestern coast of peninsular Malaysia, *A. sibogae* occurs exclusively in the uppermost reach; in contrast, *A. indicus* and *A. japonicus* uses lower streams in the same estuarine system. This may allow *A. sibogae* to avoid competition with other species for food sources and micro-habitats (see Hanamura et al., 2012).

Acetes japonicus has been recorded in both Kyushu, southern Japan and in Taiwan but not in Okinawa-jima Island, although it is located between Kyushu and Taiwan. Both *A. s. sibogae* and *A. japonicus* may occur in these regions, but we have not found any *Acetes* species other than *A. s. sibogae* on Okinawa-jima Island.

Acknowledgements

This research was supported by the JSPS KAKENHI (Grant-in-Aid for Scientific Research C) JP17K07918. This research was partially supported by Okinawa Research Core for Highly Innovative Discipline Science (ORCHIDS) project, University of the Ryukyus.

References

- Achuthankutty, C. T. & George, M. J., 1973. *Acetes sibogalis* sp. nov. (Crustacea: Decapoda, Sergestidae) from Cochin back-waters with a note on its impregnation. *Indian Journal of Marine Science*, **2**: 139-144.
- Achuthankutty, C. T. & Ayyappan Nair, S., 1976. A

- new species of sergestid shrimp, *Acetes orientalis* (Crustacea: Decapoda, Sergestidae) from Goa, central west coast of India. *Hydrobiologia*, **48**: 233-239.
- Chan, T. Y., 1998. Sergestidae. Carpenter K. E. & Niem V. H. eds. *FAO species identification guide for fishery purposes. The Living Marine Resources of the Western Central Pacific*. Vol. 2. Cephalopods, crustaceans, holothurians and sharks. FAO, Rome, 860-865 pp.
- Chihara, M. & Murano, M., 1997. *An Illustrated Guide to Marine Plankton in Japan*. Tokai University Press, Tokyo, 1574 pp. (in Japanese)
- De Grave, S., & Fransen, C. H. J. M., 2011. Carideorum catalogus: the recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). *Zoologische Mededelingen Leiden*, **85**: 195-588.
- Hanamura, Y., 1999. Occurrence of *Acetes sibogae* Hansen (Crustacea: Decapoda: Sergestidae) in Western Australia, with notes on the northern Australian population. *Records of the Western Australian Museum*, **19**: 465-468.
- Hanamura, Y., Tanaka, K., Alias M., Faizul, M. K., 2012. Ecological characteristics of hyperbenthic crustaceans in mangrove estuaries on the north-west coast of Peninsular Malaysia. *JIRCAS Working Report*, **75**: 25-34.
- Hansen, H. J., 1919. The Sergestidae of the Siboga Expedition. *Siboga-Expeditie*, **38**: 1-65, pls. 1-5.
- Hayashi, K.-I., 1992. *Dendrobranchiata crustaceans from Japanese waters*. Seibutsu Kenkyusha, Tokyo, 300 pp. (in Japanese)
- Holthuis, L. B., 1980. *FAO Species Catalogue*. Vol. 1—Shrimps and Prawns of the World: an Annotated Catalogue of Species of Interest to Fisheries. FAO Fisheries Synopsis, 125(1), Rome, 271 pp.
- Omori, M., 1975. The systematics, biogeography, and fishery of epipelagic shrimps of the genus *Acetes* (Crustacea, Decapoda, Sergestidae). *Bulletin of the Ocean Research Institute, University of Tokyo*, **7**: 1-89.
- Pérez Farfante, I. & Kensley, B., 1997. *Penaeoid and segestioid shrimps and prawns of the world*. Memoires du Museum National d'Histoire Naturelle, Paris, 175, 233 pp.
- Ravindranath, K., 1980: Shrimps of the genus *Acetes* H. Milne Edwards (Crustacea, Decapoda, Sergestidae) from the estuarine system of river Krishna. *Proceedings of the Indian Academy of Sciences (Animal Science)*, **89**: 253-273.
- Xiao, Y. & Greenwood, J. G., 1993. The biology of *Acetes* (Crustacea: Sergestidae). *Oceanography and Marine Biology: Annual Review*, **31**: 259-444.

(Received June 26, 2017; Accepted August 22, 2017)