

SCIENTIFIC REPORTS



OPEN

Atmospheric CO₂ captured by biogenic polyamines is transferred as a possible substrate to Rubisco for the carboxylation reaction

Ko Yasumoto¹, Tsuyoshi Sakata², Jun Yasumoto³, Mina Yasumoto-Hirose⁴, Shun-ichi Sato¹, Kanami Mori-Yasumoto⁵, Mitsuru Jimbo¹, Takenori Kusumi⁶ & Shugo Watabe¹

Biogenic polyamines are involved in a wide range of plant cellular processes, including cell division, morphogenesis and stress responses. However, the exact roles of biogenic polyamines are not well understood. We recently reported that biogenic polyamines that have multiple amino groups can react with CO₂ and accelerate calcium carbonate formation in seawater. The ability of biogenic polyamines to capture atmospheric CO₂ prompted us to examine their roles in photosynthesis. Here, we demonstrated that atmospheric CO₂ captured by biogenic polyamines is a candidate substrate for the carboxylation reaction of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco), which is an enzyme involved in the first major step of carbon fixation during photosynthesis, and that biogenic polyamines can accelerate the carboxylation reaction of this enzyme because of their specific affinity for CO₂. Moreover, the results of our nuclear magnetic resonance (NMR) analysis showed that putrescine, which is the most common biogenic polyamine, reacts with atmospheric CO₂ and promotes the formation of carbamate derivatives and bicarbonate in aqueous environments. A sufficient amount of CO₂ is well known to be produced by carbonic anhydrase from bicarbonate *in vivo*. The present study indicates that CO₂ would be also produced by the equilibrium reaction from carbonate produced by biogenic polyamines and would be used as a substrate of Rubisco, too. Our results may suggest a new photosynthetic research strategy that involves CO₂-concentrating mechanisms and also possibly constitutes a potential tool for reducing atmospheric CO₂ levels and, consequently, global warming.

The carboxylation reaction that fixes atmospheric CO₂ into organic compounds during photosynthesis is the first reaction of organic synthesis. This reaction, which is catalyzed by ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco), combines CO₂ with ribulose 1,5-bisphosphate (RuBP) and is the main rate-limiting reaction of photosynthesis^{1–4}. Although Rubisco is a large enzyme that has a molecular mass of approximately 550 kDa, the maximum reaction rate at 25 °C is only 15 to 30 mol CO₂ mol⁻¹ Rubisco s⁻¹⁴. Furthermore, the affinity of Rubisco for CO₂ is low: the Michaelis constant (K_m) for CO₂ at 25 °C is comparable to that of CO₂ in water equilibrated with the atmosphere^{3,4}. Therefore, plants require large amounts of Rubisco, and approximately half of leaf protein comprises this enzyme⁴.

The CO₂ used for photosynthesis in terrestrial plants diffuses from the atmosphere into the leaves through the stomata. This CO₂ then dissolves in the liquid phase of the mesophyll cell wall surface and reaches the Rubisco in the stroma of the chloroplast via the cell membrane, cytoplasm, and chloroplast envelopes⁵. This diffusion process substantially decreases the CO₂ concentration. For example, in the leaf intercellular spaces, the concentration of

¹Kitasato University School of Marine Biosciences, 1-15-1 Kitasato, Minami, Sagami-hara, Kanagawa, 252-0373, Japan. ²Biological Laboratory, Center for Natural Sciences, College of Liberal Arts and Sciences, Kitasato University, 1-15-1 Kitasato, Minami, Sagami-hara, Kanagawa, 252-0373, Japan. ³Department of Regional Agricultural Engineering, Faculty of Agriculture, University of the Ryukyus, 1 Senbaru, Nishihara, Nakagusuku, Okinawa, 903-0213, Japan. ⁴Tropical Technology Plus Co., 12-75 Suzaki, Uruma, Okinawa, 904-2234, Japan. ⁵Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University, 1314-1 Shido, Sanuki, Kagawa, 769-2193, Japan. ⁶Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1 Oookayama, Meguro-ku, Tokyo, 152-8551, Japan. Correspondence and requests for materials should be addressed to K.Y. (email: yasumoto@kitasato-u.ac.jp) or T.S. (email: sakata@kitasato-u.ac.jp)

CO₂ is reduced to 60–85% that of the atmosphere, and in the stroma, the CO₂ concentration is further reduced by 50–80%^{5–9}. Thus, the decrease in CO₂ reduces the rate of photosynthesis by approximately 2/3–1/2, and approximately half of this reduction is due to the decrease in the diffusion process from the intercellular spaces to the stroma^{10,11}. In addition to simple diffusion mentioned above, diffusion-promoting proteins such as carbonic anhydrase and aquaporins facilitate CO₂ diffusion within mesophyll cells^{12–17}. These diffusion-promoting proteins occur in the cell membrane of mesophyll cells, and aquaporin amount, which can be manipulated by altering expression levels, affects mesophyll conductance (g_m) in plants¹⁸. It is therefore necessary to accumulate more data on the characteristics and roles of the promoting mechanisms of CO₂ diffusion in leaves^{4,19,20}.

We recently reported that biogenic polyamines can capture atmospheric CO₂ and accelerate bicarbonate/carbonate formation in aqueous solutions; these findings consequently led to the formation of extracellular bacterial CaCO₃²¹. Polyamines are generally considered low-molecular-weight compounds that have multiple amino groups, are present at high concentrations in the cells of all organisms and are essential for both cell differentiation and proliferation^{22–25}. In plants, the intracellular concentrations of polyamines are a few hundred μ M to mM order²⁶. It has been reported that polyamines are localized in the vacuoles, mitochondria and chloroplasts²⁶. Moreover, chloroplasts contain a large amount of polyamines with high activities of the main polyamine biosynthetic enzymes ornithine decarboxylase (ODC) and arginine decarboxylase (ADC)²⁷. Various other functions of polyamines have been proposed, including roles as secondary messengers of plant hormones²⁶ and involvement in stress responses of plants and cyanobacteria^{28–32}. The ability of biogenic polyamines to capture atmospheric CO₂, reported by us for the first time, led us to examine the roles of biogenic polyamines in photosynthesis.

In this study, we investigated whether atmospheric CO₂ captured by biogenic polyamines could be a substrate for the carboxylation reaction of Rubisco. If Rubisco could use CO₂ incorporated in a polyamine solution as a substrate, then we could suggest an entirely new physiological function of polyamines. Therefore, we attempted to verify whether the carboxylation reaction occurs in the presence of a polyamine solution that has taken up CO₂ from the atmosphere and that serves as a substrate for commercially available, partially purified Rubisco and crude Rubisco extracted from the terrestrial plant *Fallopia japonica*. Moreover, to verify how CO₂ is incorporated from the atmosphere into the polyamine solution, we identified by nuclear magnetic resonance (NMR) the molecular species in the polyamine solution. We show that polyamines possibly contribute to CO₂ diffusion and photosynthesis. Therefore, these findings should be useful both for elucidating novel physiological functions of polyamines and for developing new methods to reduce atmospheric CO₂.

Results

Activation of Rubisco by polyamine solutions retaining CO₂. Rubisco must be activated by CO₂ and Mg²⁺ to function. Conditions consisting of 10 mM MgCl₂, 10 mM NaHCO₃, pH 7.8, and 0 °C for 10 min have been used for activation treatment³³. To verify whether polyamines supply CO₂ for Rubisco, we used polyamine solutions sufficiently incorporating CO₂ for the activation treatment of Rubisco instead of NaHCO₃ usually used as carbonate source. Figure 1 shows the comparison between Rubisco activation by piperazine and that by NaHCO₃ using commercially available Rubisco. When activation was performed using CO₂-incorporated piperazine, which is a non-natural cyclic diamine, Rubisco activity tended to increase proportionally to reaction time for 10 min. At 10 min after treatment, the activity was more than twice that at the beginning, and this time-dependent change was similar to that observed for the activation by NaHCO₃. Therefore, the activation treatments were performed using piperazine at 0 °C for 10 min.

Carboxylation reaction using CO₂-incorporated polyamine solutions as substrates. To verify whether partially purified Rubisco can utilize CO₂-incorporated polyamine solutions, we compared Rubisco activity that used different polyamines retaining CO₂ instead of NaHCO₃. The Rubisco activity was measured using a polyamine solution in which CO₂ was taken up at 20 °C for 2 days to ensure sufficient equilibrium between the CO₂ and polyamines. As shown in Fig. 2, compared with the NaHCO₃ positive control treatment, the activation treatment markedly increased the activity of Rubisco for all CO₂-incorporated polyamine solutions. The activity was 150–400% higher than that under the preactivation conditions. There were no significant differences in activity between CO₂-incorporated polyamines and NaHCO₃ [Fisher's protected least significant difference (PLSD): $P > 0.05$] prior to the activation treatment, with the exception of piperazine. In addition, after the activation treatment, the Rubisco activity from all polyamines and NaHCO₃ at the same concentrations was not significantly different (Fisher's PLSD: $P > 0.05$). Thus, CO₂-incorporated polyamine solutions were used as substrates for Rubisco.

Influence of setting time on the incorporation of CO₂ into polyamine solutions. Here, we investigated whether treatment with polyamine solutions at 20 °C for 2 days to facilitate the incorporation of CO₂ into the solutions is appropriate. For this purpose, we compared differences in the activity of partially purified Rubisco at different setting times using piperazine solutions.

As shown in Fig. 3, the activity without pretreatment for incorporation was approximately 1/5 that when NaHCO₃ was used as a substrate. The activity continued to increase from 5 to 7 hours; after 7 hours, the activity was approximately 3/4 that when NaHCO₃ was used, which was not significantly different from that when piperazine was allowed to stand for 48 hours, as shown in Fig. 3 (t-test: $P = 0.34$).

Carboxylation of crude Rubisco extracts. Similar to our examination of partially purified Rubisco, we investigated whether Rubisco in a crude extract of the leaves of the terrestrial plant *F. japonica* could be used to examine a polyamine solution incorporating CO₂ as a substrate. As shown in Fig. 4, no significant differences




Figure 1. Comparison of Rubisco activation between piperazine (Pz) solutions retaining CO₂ and a NaHCO₃ solutions. Rubisco activity was measured following activation treatment with piperazine solutions retaining CO₂ (Pz-CO₂) or NaHCO₃ solutions. The activation was carried out at 0 °C in the presence of 10 mM MgCl₂ together with either 2 mM piperazine retaining CO₂ or 10 mM NaHCO₃ (see Methods). The bars represent Rubisco activity at each time point following the activation of an enzyme solution prepared using commercially available, partially purified Rubisco. The negative control denotes the data obtained for reactions in the absence of a CO₂ supply excluding piperazine and NaHCO₃ in both the activation buffer and reaction mixture.




Figure 2. Polyamine solutions retaining CO₂ act as substrates for Rubisco. The test polyamines were dissolved in water under atmospheric conditions and then stood at 20 °C for 2 days. The solutions incorporating atmospheric CO₂ (Cad-CO₂, Put-CO₂, Spd-CO₂, Spm-CO₂ and Pz-CO₂) were used as substrates for the carboxylation reaction of partially purified Rubisco. The open columns show Rubisco activity before the activation process (nonactivated), and closed columns indicate Rubisco activity after the activation process (activated). Lower-case and upper-case letters in the figure represent the results of statistical analyses of multiple comparisons using Fisher's PLSD at the 5% significance level before and after the activation process. Bars indicate the standard error (10 mM NaHCO₃: *n* = 6, 2 mM piperazine: *n* = 10, and other polyamines: *n* = 6). The lower part of the figure shows the Rubisco activity when NaHCO₃ was used as a source of CO₂ as well as when a CO₂ source was not provided (negative control). The pH of the polyamine solutions retaining CO₂ ranged from 8.9–9.1. Abbreviations used are: Cad, cadaverine; Put, putrescine; Spd, spermidine; Pz, piperazine.

were observed between 10 mM NaHCO₃ and any of the polyamine solutions tested (Fisher's PLSD: *P* > 0.05). This trend was confirmed both before and after the activation process. In addition, following activation, the activity with polyamines as a substrate increased 10–20%, whereas that with NaHCO₃ increased 40%.




Figure 3. Influence of piperazine (Pz) solution setting time on Rubisco activity. Piperazine was dissolved in MilliQ water to a final concentration of 0.1 M and then stood at 20 °C for 48 hours to facilitate the uptake of atmospheric CO₂ (pH ranged from 8.9–9.1). To measure its activity, partially purified Rubisco was pre-activated and used as part of an enzyme solution. Bars indicate the standard error ($n = 3$).




Figure 4. Activity of a crude extract of Rubisco prepared from the leaves of *F. japonica*. A solution containing test polyamines stood at 20 °C for 2 days to facilitate the uptake of atmospheric CO₂ into the solution, after which the solution was used as a substrate for the crude enzyme extract prepared from the leaves of *F. japonica*. Refer to Fig. 2 for the measurement conditions and test methods for Rubisco activity ($n = 4$). Abbreviations used are: Cad, cadaverine; Put, putrescine; Spd, spermidine; Pz, piperazine.

Relationship between Rubisco activity and concentrations of NaHCO₃ as substrate. To investigate the relationship between the Rubisco activity and concentrations of NaHCO₃, we compared the activity at 0 to 10 mM NaHCO₃ used as substrates for the carboxylation reaction of partially purified Rubisco (Fig. S1). Rubisco activities without activation treatment were almost constant with NaHCO₃ at more than 2 mM. On the other hand, Rubisco activities after activation treatment increased accompanying increasing concentration of NaHCO₃.

Effect of CO₂-free polyamines on Rubisco activity. Solutions containing CO₂-free polyamines were used as substrates together with NaHCO₃ for the carboxylation reaction of partially purified Rubisco (Fig. S2). When the solutions containing polyamines and NaHCO₃ each of 1 mM were used as substrates, the Rubisco activities were almost equal to that of 2 mM NaHCO₃.

Enhancement of 3-phosphoglycerate production by Rubisco activity assayed in putrescine-containing medium pre-bubbled with CO₂ gas. We examined whether or not the treatment with polyamine solutions passing through CO₂ gas for 5 min was appropriately designed. CO₂ gas (99.9%) was passed through 0.1 M putrescine solution and MilliQ water (DW) at 25 °C for 0, 0.5, 1 and 5 min, respectively. These solutions containing CO₂, together with 10 mM NaHCO₃ as the positive control, were introduced




Figure 5. Enhancement of 3-phosphoglycerate production by Rubisco activity assayed in putrescine-containing medium pre-bubbled with CO₂ gas. CO₂ gas was passed through 10 mL either of 0.1 M putrescine (Put) solution or MilliQ water (DW) at room temperature for 0 to 5 min. The solutions incorporating CO₂ were used as substrates for the carboxylation of partially purified Rubisco. To activate Rubisco, either 0.5 M (final 10 mM) NaHCO₃, 0.1 M (final 2 mM) Put-CO₂ (0, 0.5, 1 and 5 min), or DW-CO₂ (0, 0.5, 1 and 5 min) was introduced to reaction solutions containing Rubisco. RuBP was added to each solution to initiate the reaction, then formic acid was added to each solution after 6 min to stop the reaction. Rubisco activity was analysed by LC-MS, detecting 3-PGA which is the direct carboxylation product of Rubisco. Different lower-case letters represent statistical significance at 5% in multiple comparisons using Tukey's test. Bars indicate the standard errors ($n = 3$).

to partially purified Rubisco, and the mixtures were allowed to stand at room temperature for 10 min to activate Rubisco. RuBP was added to the mixtures to start the carboxylation reaction, then formic acid was added to stop the reaction after 6 min. 3-phosphoglycerate (3-PGA), a direct carboxylation product by Rubisco, was subsequently measured using LC-MS. When 10 mM NaHCO₃ was used as a substrate, the concentration of 3-PGA in the resultant solution was 5.49 µg/L. Figure 5 shows the concentrations of 3-PGA after various periods in reaction as percentages of that with 10 mM NaHCO₃. The DWs which had received CO₂ gas produced 7, 19, 22 and 29% 3-PGA after 0, 0.5, 1, and 5 min, respectively. When the putrescine solution passing through CO₂ gas at a final concentration of 2 mM was used as a substrate, the concentrations of 3-PGA were 5, 44, 61 and 108% after 0, 0.5, 1, and 5 min, respectively. Thus, putrescine could capture CO₂ quickly and efficiently provide CO₂ as a substrate to Rubisco.

Results of the NMR analysis of the uptake mechanisms of putrescine and piperazine. To analyze the mechanism of CO₂ uptake by polyamines, 50 mM putrescine and piperazine were dissolved into deuterium oxide (D₂O) and then allowed to stand at 20 °C for 2–72 hours. A fixed amount of 1,4-dioxane was added as an internal standard, and the concentrations of carbamate derivatives and bicarbonate plus carbonate (HCO₃⁻ + CO₃²⁻) were calculated from ¹H-NMR and ¹³C-NMR spectra, respectively. As shown in Fig. 6, the proportion of carbamate derivatives increased beginning from the early stages. For putrescine, which is a primary diamine, the proportions of the carbamates were 2, 10, 32, 39, 53, and 55% at 2, 4, 8, 24, 48, and 72 hours, respectively. For piperazine, which is a secondary diamine, the proportions were 2, 3, 6, 19, 23, and 24% at 2, 4, 8, 24, 48, and 72 hours, respectively. In contrast, the HCO₃⁻ and CO₃²⁻ concentrations in the putrescine solution were detectable at 48 and 72 hours and were 11 and 23 mM, respectively. The HCO₃⁻ + CO₃²⁻ concentrations in the piperazine solution were detectable at 24, 48, and 72 hours and were 11, 21, and 23 mM, respectively.

NMR analysis on the rate of CO₂ uptake by putrescine in aqueous solution under 5% CO₂ condition. We analyze the rate of CO₂ uptake by putrescine in aqueous solution under 5% CO₂ condition. 50 mM putrescine in D₂O were allowed to stand for 24 hours at 25 °C in a 5% CO₂ incubator. As shown in Fig. S3, 41% of putrescine was changed to carbamate derivatives after 20 min. The carbamate derivatives increased until after 2 hours, reaching 20% after 24 hours due to the shift to cationic derivative and bicarbonate. These results indicate that putrescine had a rate of conversion to the carbamate derivative higher in 5% CO₂ than atmospheric CO₂.




Figure 6. NMR analysis results of atmospheric CO_2 incorporation by putrescine (Put) and piperazine (Pz) into aqueous solution. Changes in the relative amounts of the carbamate derivatives of amines (i.e., $-\text{CH}_2\text{NHCOO}^-$) were determined following atmospheric CO_2 uptake by putrescine and piperazine (50 mM/ D_2O), and changes in the amounts of $\text{HCO}_3^- + \text{CO}_3^{2-}$ (bicarbonate and carbonate) in the aqueous solution were also determined. Carbamate derivatives and $\text{HCO}_3^- + \text{CO}_3^{2-}$ were determined by ^1H - and ^{13}C -NMR spectra, respectively.

Discussion

Solutions containing synthetic amines, mainly alkanolamines, can capture CO_2 at high concentrations in exhaust gases³⁴. This method using artificial amines has been used since the 1930s in various industries to fix gaseous CO_2 ³⁵. In addition, another synthetic amine, polyethylene, can effectively capture CO_2 from the air³⁶. We recently showed that biogenic polyamines that are the most common biogenic amines can efficiently capture and dissolve atmospheric CO_2 in aqueous solutions at 20 to 40 °C, as in the case of artificial amines²¹. Thus, we demonstrated that biogenic polyamines play important roles in the formation of diverse calcareous skeletons in marine organisms²¹. The ability of biogenic polyamines to capture atmospheric CO_2 prompted us to examine the roles of biogenic polyamines in photosynthesis.

Our data clearly demonstrated for the first time that inorganic carbon either incorporated by polyamines from the air or derived from NaHCO_3 can act as a substrate for partially purified Rubisco (Fig. 2). Polyamines incorporating CO_2 could efficiently activate Rubisco and smoothly provide the CO_2 as a substrate to Rubisco as shown in Figs 1 and 2. These results led us to conclude that polyamines retaining CO_2 could smoothly supply this CO_2 to the amino group in the active site of Rubisco. The rate of the carboxylation reaction catalyzed by Rubisco increased with increasing setting time during which the polyamine solution took up CO_2 from the air (Fig. 3). It was also found that the substrate of this reaction is derived from atmospheric CO_2 , as the carboxylation reaction did not occur by piperazine alone (Fig. 3). This result also showed that the activation and carboxylation reaction of Rubisco is due to CO_2 supply from polyamine solution containing carbamate derivatives and carbonates, but not due to the increase of pH in the reaction solution. High Rubisco activity similar to that recorded when partially purified Rubisco was used was observed even when crude extracts of *F. japonica* were used as enzyme solutions (Fig. 4). Regardless of which polyamine- CO_2 solution was used as a substrate, the activity obtained did not significantly differ from that observed when NaHCO_3 was used. As shown in the lower part of Fig. 4, instances occurred in which moderate activity was detected even when an inorganic carbon source was not provided. This phenomenon probably occurred because the tissues of the organism contained a relatively large amount of biogenic polyamines. Also, atmospheric CO_2 is easily dissolved in the crude extract solution, even though this activity is marginal.

Figures 1 to 4 show Rubisco activities measured by the decrease in absorbance at 340 nm for NADH, accompanying the reduction of 1,3-bisphosphoglycerate (1,3-BPGA), which is a phosphorylation product of 3-PGA. Figure 5 further shows the concentrations of 3-PGA, the direct product of the carboxylation reaction of Rubisco, measured by LC-MS. Polyamines react with CO_2 in aqueous solutions, forming carbamates. The carbamate bond thus formed, though covalent, is reversible and easily releases CO_2 ³⁷. Due to such characteristics of carbamate, putrescine solution quickly captured CO_2 gas within at least 5 min, and efficiently provided the incorporated CO_2 to Rubisco as shown in Fig. 5.

The polyamines tested in this study contain different numbers of amino groups: spermine has four amino groups in one molecule, spermidine has three, and putrescine and cadaverine both have two. Piperazine is a non-natural cyclic diamine and was used for comparison. Polyamines having a larger number of amino groups incorporate more CO_2 ; however, in the experiments using partially purified and crude Rubisco extracts, no statistically significant differences among these polyamines were detected (Figs 2 and 4). Furthermore, polyamines were found to have ability of receiving CO_2 from NaHCO_3 as well as atmospheric CO_2 . Although 2 mM NaHCO_3 was not a substrate concentration that sufficiently saturates the carboxylation rate of Rubisco as shown in Fig. S1, the Rubisco activities were equal to or higher than those at 2 mM NaHCO_3 , when 1 mM NaHCO_3 plus 1 mM of polyamines were used as substrates (Fig. S2). The NMR analysis results showed that the primary diamine putrescine had a faster rate of conversion to the carbamate derivative than did the secondary diamine piperazine; after 48 hours, more than 50% of primary diamine had been converted to the carbamate derivative where the putrescine solution contained 10 mM carbonate species (Fig. 6). This phenomenon is presumably because the formation of carbamate derivatives is more likely to occur for a primary amine and because the resulting carbamate

derivative is more stable. The carbamate derivative gradually shifts to the more stable HCO_3^- . Approximately 20% of piperazine, a non-natural secondary diamine, was converted to the carbamate derivative after 72 hours, and the rate of HCO_3^- production was faster than that for putrescine. This phenomenon is probably because the carbamate derivative of the secondary amine is less stable. These results indicated that the carbamate derivatives of primary diamine biogenic polyamines may be relatively stable and more easily release CO_2 than dose bicarbonate ion only because of difference in their stability. Polyamines shown in Fig. 6 were reacted with atmospheric CO_2 in a 48-well plate with a cover, thus, the carbamate formation rate was slow because of restricted air. We reported previously that polyamines could actively capture atmospheric CO_2 and facilitate a bacterial extracellular CaCO_3 precipitation²¹. When marine bacteria were cultured on the petri dish under airtight conditions, CaCO_3 precipitation was not observed. However, once the petri dish was exposed to air, CaCO_3 was smoothly formed on the agar broth within a day. Based on these results, we concluded that polyamines facilitated the incorporation of CO_2 from the air into the culture medium²¹. We consider that the formation rate of carbamate derivatives of polyamines have the same order as Rubisco activation, because the reaction mechanisms of CO_2 absorption by polyamines are the same as that of the activation of Rubisco. Actually, when polyamines were reacted with 5% CO_2 , the rate of carbamate formation had minute order as shown in Fig. S3. There are many kinetic studies for the formation rate of carbamate derivative in synthetic amines^{34,38}, which are consistent with our results. Moreover, we roughly calculated the concentration of dissolved CO_2 in Rubisco reaction solutions. It has been reported that 0.1 M bicine (pH 8.0) containing 5 mM MgCl_2 has $\text{p}K_1 = 6.22$ ³⁹. Therefore, the concentration of dissolved CO_2 in the 10 mM NaHCO_3 solution which was used as a positive control in the present study is roughly calculated to be 105 μM by the Henderson-Hasselbalch equation. As shown in Fig. 6, the CO_2 -saturated polyamine solution contained 50% each of $[\text{HCO}_3^- + \text{CO}_3^{2-}]$ and carbamate derivatives. Therefore, the CO_2 -saturated solution in 2 mM polyamine used in the Rubisco assays is supposed to contain 1 mM HCO_3^- and dissolved CO_2 at 10.5 μM . Thus, the amount of CO_2 generated from 1 mM HCO_3^- is quite low compared with that generated from 10 mM NaHCO_3 which was used as the positive solution. Nevertheless, the Rubisco activity in the CO_2 -saturated solution in 2 mM polyamines was almost the same to or even higher than that in 10 mM NaHCO_3 (Fig. 5). Thus, the most part of dissolved CO_2 is considered to be generated from the carbamate derivatives of polyamines.

In addition, during the activation of Rubisco, the polyamine solutions retaining CO_2 showed the same effects as did NaHCO_3 solutions, which are normally used as CO_2 sources (Figs 1, 2 and 4). Rubisco is activated under weakly alkaline conditions^{40–46} and occurs via the conversion of the amino group of a lysine residue at position 201 to a carbamate derivative; it is likely that this reaction occurs only under alkaline conditions. The polyamine solution that absorbed CO_2 is weakly alkaline at pH 8.5–9.1, and many carbamate derivatives and HCO_3^- molecules exist in the solution. Therefore, it is likely that Rubisco is strongly activated, and carbonate species may be provided to serve as a substrate. However, there are several papers available, reporting that the substrate of Rubisco is CO_2 , but not HCO_3^- . For example, Cooper *et al.*⁴⁷ demonstrated that CO_2 is a better substrate for Rubisco than HCO_3^- . In the present study, we used NaHCO_3 as a carbon source for positive control in *in vitro* Rubisco assays, but not CO_2 gas, because we aimed to know the exact concentration of dissolved inorganic carbons. Under the present weakly alkaline conditions, the carboxylation reaction of Rubisco proceeded smoothly in the presence of NaHCO_3 at an mM order. This fact suggests that the sufficient amount of CO_2 was produced by the equilibrium reaction from NaHCO_3 and used as a substrate in the CO_2 fixation reaction by Rubisco. In the polyamine solution retaining carbamate and bicarbonate ion, CO_2 will be generated by the equilibrium reaction from these carbamate and bicarbonate ion. Okabe *et al.*⁴⁰ reported that hydroxylamine also enhances Rubisco activity with similar alkalisation mechanisms that we note in the presence study at CO_2 -free polyamines experiments as shown in Fig. S2. However, this substance is highly toxic and thus difficult to exist at high concentrations in intact cells⁴⁸. In contrast, polyamines are biogenic amines and exist at high concentrations within the cells. Furthermore, polyamines can directly capture atmospheric CO_2 in aqueous solution. Thus, our findings about polyamines that can play functional roles in photosynthesis are quite novel.

Our data suggest that polyamines might facilitate inorganic carbon transport from cell surfaces to Rubisco, as polyamines ensure high concentrations of retained inorganic carbon species in leaf cells. Therefore, CO_2 diffusion in leaves is the rate-limiting factor for photosynthesis in leaves^{5–9}, and it is conceivable that polyamines could contribute to photosynthesis by retaining inorganic carbon species. Furthermore, if the polyamine concentration in leaves could be increased, CO_2 could be efficiently taken up even when stomata are slightly open; this phenomenon may help prevent moisture loss from the stomata during desiccation and suggests that polyamines may also be involved in drought tolerance of plants³². As mentioned before, chloroplasts contain a large amount of polyamines with ODC and ADC, both of which are main polyamine biosynthetic enzymes^{26,27}. Based on our results, we speculate the role of polyamines in inorganic carbons concentration mechanisms in plants as follows. The high concentrations of polyamines in the cytosol and chloroplasts provide high concentration of intracellular inorganic carbons, where CO_2 generated by the equilibrium reaction from these inorganic carbons (carbamates) and bicarbonate ion will be used as a substrate of Rubisco. Furthermore, a part of the polyamines synthesized in the cytosol permeate the intercellular air spaces through polyamine transporters⁴⁹. These polyamines could capture CO_2 contained in intercellular air spaces and transport to intercellular fluid, then contribute to CO_2 diffusion. Of course, a large amount of CO_2 is well known to be produced by carbonic anhydrase from bicarbonate and used as a substrate of Rubisco *in vivo*.

Moreover, the ability of polyamines to produce bicarbonate-carbonate in aqueous solution decreases at low temperatures because this ability requires high temperatures for the efficient hydration reaction of the carbamate derivatives to bicarbonate²¹. This property of polyamines might be related to the promotion of polyamine biosynthesis during exposure to low temperatures²⁸. This novel mechanism for CO_2 fixation by Rubisco involving biogenic polyamines provides a new strategy for photosynthetic research and suggests a new CO_2 -removal concept that could reduce both atmospheric CO_2 levels and global warming.

Methods

Preparation of polyamine solutions for Rubisco assay. To incorporate atmospheric CO₂ into polyamine solutions, solutions (2 ml) containing piperazine (Wako Pure Chemicals, Osaka, Japan) and biogenic polyamines (putrescine, spermidine, cadaverine and spermine; Wako Pure Chemicals, Osaka, Japan) each at 0.1 M were added to multidishes (24 wells, diameter of 15 mm), which stood for 48 hours at 20 °C. The resultant polyamine solutions at pH 8.9–9.1 were used as carbonate sources in Rubisco assays.

To activate Rubisco, the enzyme was preincubated in the presence of the polyamine solutions retaining CO₂ instead of NaHCO₃ solutions³³.

To investigate the time-dependent changes in Rubisco activity, solutions (2 ml) containing 0.1 M piperazine were added to multidishes (24 wells, diameter of 15 mm), which stood for 0, 5, 7 and 48 hours at 20 °C. The resultant piperazine solutions at pH 8.9–9.1 were used as carbonate sources in Rubisco assays. Ten millimolar solutions of NaHCO₃ served as positive controls.

Rubisco assays. Partially purified Rubisco (0.05 units mg⁻¹ solid; Sigma-Aldrich, St. Louis, MO, USA) and crude extracts from the leaves of *F. japonica* Houtt. var. *japonica* were used in Rubisco assays.

The partially purified Rubisco was dissolved in buffer [100 mM hydroxyethyl piperazineethanesulfonic acid (HEPES), 10 mM dithiothreitol (DTT), 5 mM MgCl₂ and 1 mM EDTA, pH 7.8] at 10 m units 75 μl⁻¹ as an enzyme solution. The buffer was equilibrated with pure N₂ gas in order to exclude CO₂.

The leaves of *F. japonica* were collected on 25 September 2012 at Kitasato University (Sagamihara, Kanagawa, Japan). Two leaf discs (0.49 cm² each) were punched, frozen immediately in liquid N₂, and then maintained at -80 °C until assays. Two frozen leaf discs were rapidly homogenized in a chilled mortar filled with 1 ml of CO₂-free extraction buffer [100 mM HEPES, 10 mM DTT, 5 mM MgCl₂, 1 mM EDTA, 2% (w/v) PVP40, 1% (v/v) Triton X-100 and 0.2 mM leupeptin, pH 7.8] for 3 min. The extraction buffer was equilibrated with pure N₂ gas in order to exclude CO₂ prior to the extraction. The homogenate was centrifuged at 17400 × g for 2 min at 4 °C, after which the supernatant was used immediately to assay the initial activity of Rubisco (nonactivated).

The total activity of Rubisco (activated) was also determined following the activation of a 160 μl enzyme solution and was achieved by preincubation for more than 10 min at 0 °C in 40 μl of the activation buffer (75 mM HEPES, 10 mM MgCl₂ and 10 mM NaHCO₃ or 2 mM polyamine solution).

Rubisco activity was determined in accordance with the spectrophotometric method of Lilley and Walker⁵⁰, which was partly modified by Sakata *et al.*⁵¹. The reaction mixture (100 mM bicine buffer containing 5 mM MgCl₂, 5 mM creatine phosphate, 1 mM ATP, 5 units ml⁻¹ creatine kinase, 5 units ml⁻¹ 3-phosphoglycerate kinase, 5 units of glyceraldehyde-3-phosphate dehydrogenase and 0.1 mM NADH, pH 8.2) was treated with pure N₂ gas for 30 min in order to exclude CO₂. All chemicals and enzymes were commercially available. A 2925 μl reaction mixture that contained 150 μl of 6 mM RuBP and either 60 μl of 0.5 M NaHCO₃ or 0.1 M polyamine solutions was added to a cuvette under a N₂ gas atmosphere at 25 °C. After confirming no more decrease in the absorbance at 340 nm, the Rubisco activity was measured by the addition of 75 μl of partially purified Rubisco or the crude extract to the cuvette. Approximately 5 min elapsed between the start of homogenization and the start of assay. Rubisco activity was recorded by the decrease in absorbance at 340 nm and was corrected by a blank assay, in which the reaction mixture did not contain RuBP. To determine Rubisco activity per unit protein, the protein content in the extract was assayed according to the method of Bradford⁵² using a protein assay kit (BIO-RAD, Hercules, CA, USA).

Rubisco activity in putrescine solution passing through CO₂ gas as detected with the increased concentrations of 3-PGA by LC-MS. CO₂ gas (99.9%) was passed through 10 mL either of 0.1 M putrescine solution or MilliQ water, which had been pre-treated with N₂ gas to exclude CO₂, for 0, 0.5, 1 and 5 min at room temperature. The solutions incorporating CO₂ were used as substrates for the carboxylation reaction of partially purified Rubisco. A 905 μl of CO₂-free reaction mixture (0.1 M bicine, pH 8.2, containing 5 mM MgCl₂), 25 μl of the enzyme solution of Rubisco, and either 20 μl of 0.5 M NaHCO₃, 0.1 M putrescine solutions incorporating CO₂, or MilliQ water were introduced to 1.5 mL tube. The mixed solutions were allowed to stand for 10 min at 25 °C to activate Rubisco. Then, each of the solutions were added with 50 μl of 6 mM RuBP to start the reaction, and after 6 min later with 200 μl of formic acid to stop the reaction. 3-phosphoglyceric acid (3-PGA), the direct product of the carboxylation reaction of Rubisco was measured using LC-MS. In brief, high-resolution hybrid quadrupole-time-of-flight mass spectrometer (Triple TOF 5600+, SCIEX) operated in a negative ion mode was coupled to reversed phase chromatography via electrospray ionization and scanned from *m/z* 50 to 600 at high resolution. LC separation was achieved on an InertSustain C18 column (2.1 mm × 150 mm, 3 μm particle size, GL Sciences) using a gradient of solvent A (10 mM tributylamine + 10 mM acetic acid in water) and solvent B (methanol). The gradient was: 0 min, 0% B; 1 min, 0% B; 1.5 min, 15% B; 3 min, 15% B; 8 min, 50% B; 10 min, 100% B; 11 min, 100% B; 11.5 min, 0% B; 17 min, 0% B at a flow rate of 200 μl min⁻¹. Quantification of 3-PGA for its deprotonated molecule [M-H]⁻ at *m/z* 184.9857 was performed using MutliQuant integration software (SCIEX). The standard compound of 3-PGA was purchased from Sigma-Aldrich (USA).

LC separation was achieved on an InertSustain C18 column (2.1 mm × 150 mm, 3 μm particle size, GL Sciences) using a gradient of solvent A (10 mM tributylamine + 10 mM acetic acid in water) and solvent B (methanol). The gradient was: 0 min, 0% B; 1 min, 0% B; 1.5 min, 15% B; 3 min, 15% B; 8 min, 50% B; 10 min, 100% B; 11 min, 100% B; 11.5 min, 0% B; 17 min, 0% B at a flow rate of 200 μl min⁻¹. Quantification of 3-PGA for its deprotonated molecule [M-H]⁻ at *m/z* 184.9857 was performed using MutliQuant integration software (SCIEX). The standard compound of 3-PGA was purchased from Sigma-Aldrich (USA).

NMR analysis of the uptake mechanisms of putrescine and piperazine. Two-milliliter solutions of D₂O containing 50 mM putrescine or 50 mM piperazine were added to multidishes (24 wells, diameter of 10 mm). The multidishes stood at 20 °C for 2, 4, 8, 24, 48, or 72 hours. The resultant carbamate and HCO₃⁻ + CO₃²⁻ (bicarbonate and carbonate) were characterized by using ¹H-NMR and ¹³C-NMR^{21,53-55}. An internal standard of 1,4-dioxane (5 μl, diluted 10-fold with D₂O) was added to each NMR sample. The yield of carbamate derivatives was estimated by the ¹H-NMR spectrum based on the area ratio of integration beneath the peaks for the α-methylene proton of the amines and their carbamates^{21,53}. To determine the concentrations of HCO₃⁻ + CO₃²⁻, solutions of D₂O containing 5 to 50 mM NaHCO₃ were measured with ¹³C-NMR^{54,55}. A calibration curve was

then obtained from the area ratio of integration beneath the peaks for both dioxane and $\text{HCO}_3^- + \text{CO}_3^{2-}$ from the spectra of the NaHCO_3 solutions, and the concentration was estimated by the ratio of dioxane to $\text{HCO}_3^- + \text{CO}_3^{2-}$ in each sample. The ^1H - and ^{13}C -NMR spectra were recorded using a Bruker AVANCE (600 and 700 MHz) spectrometer.

NMR analysis on CO_2 incorporation by putrescine in aqueous solution under 5% CO_2 condition. To examine the reactivity of putrescine under 5% CO_2 condition, one-milliliter solutions of D_2O containing 50 mM putrescine were introduced into multidishes (24 wells, diameter of 10 mm). The multidishes stood at 25 °C for 24 hours in a 5% CO_2 incubator. The resultant carbamates were characterized by using ^1H -NMR as shown in Fig. S3.

References

- Farquhar, G. D. & Von Caemmerer, S. Modelling of photosynthetic response to environmental conditions. In *Physiological Plant Ecology II* (pp. 549–587). Springer Berlin Heidelberg (1982).
- Evans, J. R. & Seemann, J. R. The allocation of protein nitrogen in the photosynthetic apparatus: costs, consequences, and control. *Photosynthesis*, 183–205 (1989).
- Andrews, T. J. & Whitney, S. M. Manipulating ribulose biphosphate carboxylase/oxygenase in the chloroplasts of higher plants. *Archives of Biochemistry and Biophysics* **414**, 159–169 (2003).
- Terashima, I., Hanba, Y. T., Tholen, D. & Niinemets, Ü. Leaf functional anatomy in relation to photosynthesis. *Plant Physiology* **155**, 108–116 (2011).
- Evans, J. R. & Loreto, F. Acquisition and diffusion of CO_2 in higher plant leaves. In *Photosynthesis* (pp. 321–351). Springer Netherlands (2000).
- Terashima, I., Hanba, Y. T., Tazoe, Y., Vyas, P. & Yano, S. Irradiance and phenotype: comparative eco-development of sun and shade leaves in relation to photosynthetic CO_2 diffusion. *Journal of Experimental Botany* **57**, 343–354 (2005).
- Flexas, J., Ribas-Carbó, M., Diaz-Espejo, A., Galmés, J. & Medrano, H. Mesophyll conductance to CO_2 : current knowledge and future prospects. *Plant, Cell & Environment* **31**, 602–621 (2008).
- Evans, J. R. Potential errors in electron transport rates calculated from chlorophyll fluorescence as revealed by a multilayer leaf model. *Plant and Cell Physiology* **50**, 698–706 (2009).
- Niinemets, Ü., Diaz-Espejo, A., Flexas, J., Galmés, J. & Warren, C. R. Importance of mesophyll diffusion conductance in estimation of plant photosynthesis in the field. *Journal of Experimental Botany* **60**, 2271–2282 (2009).
- Sakata, T. & Yokoi, Y. Analysis of the O_2 dependency in leaf-level photosynthesis of two *Reynoutria japonica* populations growing at different altitudes. *Plant, Cell & Environment* **25**, 65–74 (2002).
- Yamori, W., Suzuki, K., Noguchi, K. O., Nakai, M. & Terashima, I. Effects of Rubisco kinetics and Rubisco activation state on the temperature dependence of the photosynthetic rate in spinach leaves from contrasting growth temperatures. *Plant, Cell & Environment* **29**, 1659–1670 (2006).
- Yagawa, Y., Shiraiwa, Y. & Miyachi, S. Carbonic anhydrase from the blue-green alga (cyanobacterium) *Anabaena variabilis*. *Plant and Cell Physiology* **25**(5), 775–783 (1984).
- Tsuzuki, M. & Miyachi, S. The function of carbonic anhydrase in aquatic photosynthesis. *Aquatic Botany* **34**(1–3), 85–104 (1989).
- Terashima, I. & Ono, K. Effects of HgCl_2 on CO_2 dependence of leaf photosynthesis: evidence indicating involvement of aquaporins in CO_2 diffusion across the plasma membrane. *Plant and Cell Physiology* **43**, 70–78 (2002).
- Hanba, Y. T. *et al.* Overexpression of the barley aquaporin HvPIP2; 1 increases internal CO_2 conductance and CO_2 assimilation in the leaves of transgenic rice plants. *Plant and Cell Physiology* **45**, 521–529 (2004).
- Giordano, M., Beardall, J. & Raven, J. A. CO_2 concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annual Review of Plant Biology* **56**, 99–131 (2005).
- Flexas, J. *et al.* Tobacco aquaporin NtAQP1 is involved in mesophyll conductance to CO_2 *in vivo*. *The Plant Journal* **48**, 427–439 (2006).
- Warren, C. R. & Dreyer, E. Temperature response of photosynthesis and internal conductance to CO_2 : results from two independent approaches. *Journal of Experimental Botany* **57**, 3057–3067 (2006).
- Sakata, T., Nakano, T. & Kachi, N. Effects of internal conductance and Rubisco on the optimum temperature for leaf photosynthesis in *Fallopia japonica* growing at different altitudes. *Ecological Research* **30**, 163–171 (2015).
- Tazoe, Y. & Santrucek, J. Superimposed behaviour of g_m under ABA-induced stomata closing and low CO_2 . *Plant, Cell & Environment* **38**, 385–387 (2015).
- Yasumoto, K. *et al.* Biogenic polyamines capture CO_2 and accelerate extracellular bacterial CaCO_3 formation. *Marine Biotechnology* **16**, 465–474 (2014).
- Tabor, C. W. & Tabor, H. Polyamines in microorganisms. *Microbiological Reviews* **49**, 81 (1985).
- Guergué, A., Claparols, I., Santos, M. & Torné, J. M. Modulator effect of DL- α -difluoromethylarginine treatments on differentiation processes of young maize calluses. *Plant Growth Regulation* **21**, 7–14 (1997).
- Kakkar, R. K., Nagar, P. K., Ahuja, P. S. & Rai, V. K. Polyamines and plant morphogenesis. *Biologia Plantarum* **3**, 1–11 (2000).
- Kashiwagi, K. & Igarashi, K. Identification and assays of polyamine transport systems in *Escherichia coli* and *Saccharomyces cerevisiae*. *Polyamines: Methods and Protocols*, 295–308 (2011).
- Slocum, R. D. & Flores, H. E. *Biochemistry and Physiology of Polyamines in Plants*. CRC press (1991).
- Andreadakis, A. & Kotzabasis, K. Changes in the biosynthesis and catabolism of polyamines in isolated plastids during chloroplast photodevelopment. *Journal of Photochemistry and Photobiology B: Biology* **33**, 163–170 (1996).
- Bouchereau, A., Aziz, A., Larher, F. & Martin-Tanguy, J. Polyamines and environmental challenges: recent development. *Plant Science* **140**, 103–125 (1999).
- Velikova, V., Yordanov, I. & Edreva, A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Science* **151**, 59–66 (2000).
- He, L., Nada, K., Kasukabe, Y. & Tachibana, S. Enhanced susceptibility of photosynthesis to low-temperature photoinhibition due to interruption of chill-induced increase of S-adenosylmethionine decarboxylase activity in leaves of spinach (*Spinacia oleracea* L.). *Plant and Cell Physiology* **43**, 196–206 (2002).
- Incharoensakdi, A., Jantaro, S., Raksajit, W., & Mäenpää, P. Polyamines in cyanobacteria: biosynthesis, transport and abiotic stress. *Current research, technology and education topics in applied microbiology and microbial biotechnology. Formatex, Badajoz*, 23–32 (2010).
- Sequera-Mutiozabal, M., Tiburcio, A. F. & Alcázar, R. Drought stress tolerance in relation to polyamine metabolism in plants. *Drought Stress Tolerance in Plants, Springer International Publishing* **1**, 267–286 (2016).
- Sakata, T., Kachi, N. & Yokoi, Y. Quantitative evaluation of the counterbalance between photosynthetic stimulation and depression caused by low partial pressure of O_2 and CO_2 in alpine atmospheres. *Polar Science* **1**, 55–62 (2007).
- Conway, W. *et al.* Toward rational design of amine solutions for PCC applications: the kinetics of the reaction of CO_2 (aq) with cyclic and secondary amines in aqueous solution. *Environmental Science & Technology* **46**, 7422–7429 (2012).

35. Rochelle, G. T. Amine scrubbing for CO₂ capture. *Science* **325**, 1652–1654 (2009).
36. Goeppert, A. *et al.* Carbon dioxide capture from the air using a polyamine based regenerable solid adsorbent. *Journal of the American Chemical Society* **133**, 20164–20167 (2011).
37. Conway, W. *et al.* Reactions of CO₂ with aqueous piperazine solutions: formation and decomposition of mono- and dicarbamic acids/carbamates of piperazine at 25.0 °C. *The Journal of Physical Chemistry A* **117**, 806–813 (2013).
38. Vaidya, P. D. & Kenig, E. Y. CO₂-Alkanolamine Reaction Kinetics: A review of recent studies. *Chemical Engineering & Technology* **30**, 1467–1474 (2007).
39. Makino, A., Mae, T. & Ohira, K. Enzymic properties of ribulose-1, 5-bisphosphate carboxylase/oxygenase purified from rice leaves. *Plant Physiology* **79**, 57–61 (1985).
40. Okabe, K. I., Codd, G. A. & Stewart, W. D. Hydroxylamine stimulates carboxylase activity and inhibits oxygenase activity of cyanobacterial RuBP carboxylase/oxygenase. *Nature* **279**, 525–527 (1979).
41. McCurry, S. D., Pierce, J., Tolbert, N. E. & Orme-Johnson, W. H. On the mechanism of effector-mediated activation of ribulose bisphosphate carboxylase/oxygenase. *Journal of Biological Chemistry* **256**, 6623–6628 (1981).
42. Badger, M. R. & Lorimer, G. H. Interaction of sugar phosphates with the catalytic site of ribulose-1, 5-bisphosphate carboxylase. *Biochemistry* **20**, 2219–2225 (1981).
43. Mizioro, H. M. & Lorimer, G. H. Ribulose-1, 5-bisphosphate carboxylase-oxygenase. *Annual Review of Biochemistry* **52**, 507–535 (1983).
44. Lilley, R. M. & Portis, A. R. Activation of ribulose-1, 5-bisphosphate carboxylase/oxygenase (Rubisco) by rubisco activase. *Plant Physiology* **94**, 245–250 (1990).
45. Majeau, N. & Coleman, J. R. Correlation of carbonic anhydrase and ribulose-1, 5-bisphosphate carboxylase/oxygenase expression in pea. *Plant physiology* **104**, 1393–1399 (1994).
46. Portis, A. R. & Parry, M. A. Discoveries in Rubisco (ribulose 1, 5-bisphosphate carboxylase/oxygenase): a historical perspective. *Photosynthesis Research* **94**, 121–143 (2007).
47. Cooper, T. G., Filmer, D., Wishnick, M. & Lane, M. D. The active species of “CO₂” utilized by ribulose diphosphate carboxylase. *Journal of Biological Chemistry* **244**, 1081–1083 (1969).
48. Evelo, C. T., Spooren, A. A., Bisschops, R. A., Baars, L. G. & Neis, J. M. Two mechanisms for toxic effects of hydroxylamines in human erythrocytes: involvement of free radicals and risk of potentiation. *Blood cells, Molecules, and Diseases* **24**, 280–295 (1998).
49. Fujita, M. *et al.* Natural variation in a polyamine transporter determines paraquat tolerance in Arabidopsis. *Proceedings of the National Academy of Sciences* **109**, 6343–6347 (2012).
50. Lilley, R. M. & Walker, D. A. An improved spectrophotometric assay for ribulosebisphosphate carboxylase. *Biochimica et Biophysica Acta (BBA)-Enzymology* **358**, 226–229 (1974).
51. Sakata, T., Nakano, T., Iino, T. & Yokoi, Y. Contrastive seasonal changes in ecophysiological traits of leaves of two perennial Polygonaceae herb species differing in leaf longevity and altitudinal distribution. *Ecological Research* **21**, 633–640 (2006).
52. Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**, 248–254 (1976).
53. Ume, C. S. & ALPER, E. Reaction kinetics of carbon dioxide with 2-amino-2-hydroxymethyl-1, 3-propanediol in aqueous solution obtained from the stopped flow method. *Turkish Journal of Chemistry* **36**, 427–435 (2012).
54. Barzagli, F., Mani, F. & Peruzzini, M. A ¹³C NMR study of the carbon dioxide absorption and desorption equilibria by aqueous 2-aminoethanol and N-methyl-substituted 2-aminoethanol. *Energy & Environmental Science* **2**, 322–330 (2009).
55. Mani, F., Peruzzini, M. & Stoppioni, P. CO₂ absorption by aqueous NH₃ solutions: speciation of ammonium carbamate, bicarbonate and carbonate by a ¹³C NMR study. *Green Chemistry* **8**, 995–1000 (2006).

Acknowledgements

This work was supported partly by grants-in-aid from the Japan Society for the Promotion of Science (KAKENHI Grant Numbers 25850145, 15K07587, 16K07526 and 16K00518), the Institute for Fermentation Osaka, and Kitasato University for Young Researchers.

Author Contributions

K.Y., T.S., J.Y. and M.Y.H. designed the study. Most experiments were performed by K.Y., T.S. and S.S.; in collaboration with S.W., K.Y. and T.S. wrote the paper. Rubisco experiments were performed by T.S. and S.S. and NMR spectroscopy was performed by K.Y., K.M.Y. and K.T. All authors discussed the results and commented on the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-35641-8>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018