

琉球大学学術リポジトリ

Upper Lip Reconstruction after Oncologic Resection by a Sliding Advancement Cheek Flap with Buccal Mucosal Eversion

メタデータ	言語: 出版者: Lippincott, Williams & Wilkins 公開日: 2020-06-25 キーワード (Ja): キーワード (En): 作成者: Nishihara, Kazuhide, Nakasone, T., Matayoshi, A., Maruyama, T., Arasaki, A. メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/46287

Upper Lip Reconstruction after Oncologic Resection by a Sliding Advancement Cheek Flap with Buccal Mucosal Eversion

Kazuhide Nishihara, DDS, PhD*†; T. Nakasone, DDS, PhD†; A. Matayoshi, DDS, PhD†; T. Maruyama, DDS†; A. Arasaki, DDS, PhD*†

Sir:

The objective of lip reconstruction is to achieve both functional and esthetic competence in mastication, phonation, facial expression, and facial balance. Upper lip defects can often be reconstructed using an advancement/rotation flap with Burrow's triangle excision and a crosslip flap such as the Abbe flap or Estlander flap.¹⁻⁵ Martin et al³ reported that plastic surgeons' first choice

for reconstruction of upper lip defects was the cheek advancement flap, followed by the Abbe flap. Excellent esthetic and functional consequences for the reconstruction of large upper lip defects have yet to be established.

A 71-year-old man presented with a 5-month history of an ulcer in the right upper lip. The patient had visited a dental clinic and was clinically diagnosed with a decubital ulcer caused by contact between the canine




Fig. 1. Squamous cell carcinoma of the upper lip reconstructed with an advancement cheek flap. Preoperatively planned marking for bilateral cheek advancement with Burrow's triangle of the nasal ala and lower lip (removed at a, b, c, and d regions) for a large defect after resection of the upper lip carcinoma. The upper lip reconstruction was performed using the 2 sliding rotations and an advancement cheek mucosal flap (A and B flaps). The upper vermilion reconstruction was accomplished with buccal mucosal eversion (red arrow regions).

From the *Department of Oral and Maxillofacial Functional Rehabilitation, Graduate School of Medicine, University of the Ryukyus, Nakagami, Okinawa, Japan; and †Department of Oral and Maxillofacial Surgery, University of the Ryukyus Hospital, Nakagami, Okinawa, Japan.

Copyright © 2016 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. All rights reserved. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Plast Reconstr Surg Glob Open 2016;4:e1100 doi:10.1097/GOX.0000000000001100; Published online 10 November 2016.

teeth and right upper lip. Clinical examination at our institution revealed a 33×17-mm wedge-shaped ulcer with surrounding indurations involving the right upper lip region. The patient underwent a biopsy of the lesion under local anesthesia to achieve a pathological diagnosis, and histopathological examination revealed moderately differentiated squamous cell carcinoma. The patient underwent excision of the tumor mass, total neck dissection, and reconstruction of the right upper lip under general anesthesia.

The upper lip reconstruction was performed using 2 sliding advancement cheek flaps and buccal mucosal eversion. A curvilinear region involving Burrow's triangle was designed



Fig. 2. One-year postoperative view. Note the presence of the upper vermilion recruited by buccal mucosal eversion. The patient was able to close the upper and lower lips and masticate using dentures.

along the bilateral alar base of the nose and cheek skin crease to allow for sliding of the advancement cheek flap. The length of the lower incision line was planned to be equal to that of the horizontal defect. Next, wide full-thickness resection was performed to a horizontal diameter of approximately 50mm around the malignant tumor and involved almost three-fourths of the upper lip. After resection of the triangular curvilinear region, the medial edges of each flap were inset with 3-layer closure to provide adequate oral sphincter function. The lower resections were performed through the subcutaneous tissue and muscle to the level of the mucosa. An inferior-based buccal mucosal flap was reversed at the anterior aspect of this incision onto the lower triangular region to create the lateral commissure and upper red lip. Upper vermilion reconstruction was accomplished with buccal mucosal eversion of the lower triangular region (Fig. 1).

The Abbe flap involves a 2-stage reconstruction with risk of relative microstomia, lower lip distortion, and

vascular robustness when applied to large defects.^{3,4} However, our single-stage procedure avoids the development of microstomia and allows for closure of the upper and lower lips and mastication using dentures (Fig. 2). Our surgical technique provides reliable reconstruction of large upper lip defects reaching on from the commissure to the nasal sill and the ala base. We recommend this surgical technique for repair of large defects measuring more than three-fourths of the upper lip.

Kazuhide Nishihara, DDS

Department of Oral and Maxillofacial Functional Rehabilitation
Graduate School of Medicine
University of the Ryukyus
207 Uehara Nishihara
Nakagami, Okinawa 903-0215, Japan
E-mail: kazuhide@med.u-ryukyu.ac.jp

DISCLOSURE

The authors have no financial interest to declare in relation to the content of this article. The Article Processing Charge was paid for by the authors.

REFERENCES

1. Williams EF 3rd, Setzen G, Mulvaney MJ. Modified Bernard-Burow cheek advancement and cross-lip flap for total lip reconstruction. *Arch Otolaryngol Head Neck Surg.* 1996;122:1253–1258.
2. Anvar BA, Evans BC, Evans GR. Lip reconstruction. *Plast Reconstr Surg.* 2007;120:57e–64e.
3. Martin TJ, Zhang Y, Rhee JS. Options for upper lip reconstruction: a survey-based analysis. *Dermatol Surg.* 2008;34:1652–1658.
4. Baumann D, Robb G. Lip reconstruction. *Semin Plast Surg.* 2008;22:269–280.
5. García de Marcos JA, Heras Rincón I, González Córcoles C, et al. Bilateral reverse Yu flap for upper lip reconstruction after oncologic resection. *Dermatol Surg.* 2014;40:193–196.