

琉球大学学術リポジトリ

Growth Parameters and Yield Response of Maize to Water Stress and Nitrogenous Fertilizer

メタデータ	言語: 出版者: 沖縄農業研究会 公開日: 2009-01-29 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/0002015513

Growth Parameters and Yield Response of Maize to Water Stress and Nitrogenous Fertilizer

M.S.U.Talukder¹, S.M. Shirazi², M.A.Hossain³, H. Dey¹, M.A. Hye⁴

INTRODUCTION

In Bangladesh, maize is the third most important cereal crop and covers 2834 hectares of land with an annual production of 3000 tons (BBS,1997). Proper growth and development of maize needs favourable soil moisture in the root zone. The moisture content in the soil gradually decreases with the passing of time during dry season. Limited water supply during the growing season results in soil and plant water deficits and reduces maize yields (Gordon *et al.*, 1995). Proper time and supplemental irrigation should be realized in irrigation scheduling for the most effective use of available water in optimizing maize production. Water deficit has little effect on timing of emergence, number of leaves per plant but delayed tasseling initiation and silking, reduced plant height and vegetative growth of maize (Abrecht and Carberry, 1993). Heading to milking stage is the most sensitive period of water stress and has ultimate impact on grain yield (Shaozhong and Minggang, 1992). Improper scheduling of irrigation results not only in wastage of

water but also decrease the crop growth and yield.

Nitrogen for maize cultivation is equally important to realize the yield potential (Talukder, 1985). Among different elements of Bangladesh soil, nitrogen is the key input for achieving higher yield of maize. But nitrogenous fertilizer may be increased to a certain level and thereafter it has got adverse effect (Gupta and Gautam; 1994, Singh *et al.*, 1996). Irrigation water dissolved the fertilizers and made available to the crop for proper growth and development. Therefore, an attempt has been made to evaluate the effect of irrigation and nitrogen on the performance of maize.

MATERIALS AND METHOD

The experiment was conducted in the Agronomy Research Field of Bangladesh Agricultural University, Mymensingh during November 1998 to April 1999 to evaluate the response of maize (cv. Bornali) to water stress and nitrogenous fertilizer. The experiment included two factors, namely i) five irrigation regimes with IW/CPE ratios of 0.0, 0.2, 0.5, 0.8, and 1.0 and was applied at 37, 58 and 75 days after sowing (DAS) and ii) four nitrogen doses i.e. 00, 70, 100 and 120 N kg/ha. The experiment was laid out in a split plot design with 3 replications assigning 5 irrigation treatments to main plots and 4 fertility treatments to sub-plots at random. Texturally, the soil was silt loam. The land was prepared by the power tiller. Seeds were sown on 19 November 1998 by dropping seeds by hand with 70 cm x 25 cm spacing. The unit plot size was 4 m x 1.5 m (5

¹ Department of Irrigation and Water Management, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

² Studies on Integrated Farming, Bangladesh Agricultural University, Mymensingh - 2202, Bangladesh.

³ Laboratory of Irrigation and Water Management, Faculty of Agriculture, University of the Ryukyus, Okinawa 903-0213, Japan.

⁴ Department of Mathematics, Agricultural University, Mymensingh-2202, Bangladesh.

sq.m). Triple superphosphate (TSP) and muriate of potash (MP) were applied at the rate of 100 Kg P₂O₅ and 80 Kg K₂O/ha, respectively (BARC, 1997). One third of the nitrogen along with whole TSP and MP were applied at the time of final land preparation. The rest two-third urea was top dressed in equal two splits at 35 and 65 days after sowing. The maize was harvested on 6 April 1999. Intercultural operations were made as and when necessary to keep the crop free from weeds and to protect from diseases. Soil moisture was determined at 34, 39, 54, 60, 73 and 77 DAS from each main plot. Soil samples were also collected from unit plots during land preparation and at harvest to determine the physico-chemical properties of soil. Plant height, root length, ear length, ear breath, ear/plant, kernel/ear, 100 kernel weight, grain and straw yields were recorded. Data were analyzed following analysis of variance technique and mean difference were adjudged by Duncan's Multiple Range Test (Gomez and Gomez, 1984)

RESULTS AND DISCUSSION

Effect of irrigation regimes on the yield and yield contributing characters of maize are presented in Table 1. The yield and yield parameters were significantly affected due to application of irrigation water. The highest plant height (273.50 cm) was observed with IW/CPE ratio of 0.8 irrigation treatment and the lowest (249.80 cm) in control. Availability of well distributed soil moisture at different growth stages due to irrigation enhanced the growth of plant. Similar effect of irrigation on plant height was also reported by Gordon *et al.*, (1995) and Ne Smith and Ritchie (1992). Root lengths were significantly increased with the application of irrigation water. The highest root length (29.50 cm) was produced in IW/CPE ratio of 0.2 and 0.8 irrigation treatments and was statistically similar with other irrigation treatments except control (26.42 cm). Similar result was found by Dai *et al.*, (1990). Due to application of irrigation water ear length was significantly increased (Table 1). The highest ear length (19.72 cm) was produced by IW/CPE ratio of 0.2 irrigation treatment and was statistically similar with

Table 1. Effect of irrigation regimes on the yield and yield contributing characters of maize.

IW/CPE ratio	Plant height (cm)	Root length (cm)	Ear length (cm)	Ear breath (cm)	Ears/plant (no.)	Kernels/Ear (no.)	100 Kernel weight (g)	Grain yield (t/ha)	Stover yield (t/ha)
0	249.80c	26.42b	18.46b	4.04b	1.12	310.30d	24.40d	3.85d	6.84d
0.2	265.90ab	29.50a	19.72a	4.88a	1.35	372.50a	26.69c	5.77b	9.53b
0.5	271.50a	28.87a	20.21a	4.78a	1.38	351.00b	31.75a	6.77a	11.13a
0.8	273.50a	29.50a	18.34b	4.23b	1.38	333.10bc	31.05b	5.61b	8.35c
1.0	256.80bc	29.38a	18.25b	4.30b	1.20	327.10cd	27.03c	4.80c	7.87c
S _x	2.05	0.17	0.11	0.06	-	4.34	0.12	0.10	0.11
Level of significance	0.01	0.01	0.01	0.01	NS	0.01	0.01	0.01	0.01

Note : Figure in a column having common letter(s) do not differ significantly but dissimilar letter differ significantly; NS = Not significant.

the ratio of 0.5 irrigation treatment. The lowest ear length (18.25 cm) was obtained with IW/CPE ratio of 1.0 irrigation treatment and was statistically identical with 0.8 and control. Ear breath was significantly affected by irrigation water and followed similar trend as in ear length (Table 1). Number of kernels / ear were significantly affected due to application of irrigation water. IW/CPE ratio of 0.2 irrigation treatment produced the highest kernel number/ear (372.50) and the lowest (310.30) in control. A significant variation was recorded for 100 kernel weight owing to differences in irrigation treatments (Table1). Influence of irrigation on grain yield was statistically significant. The highest grain yield (6.77 t/ha) was obtained with IW/CPE ratio of 0.5 irrigation treatment and the lowest (4.80 t/ha) under no stress condition (IW/CPE = 1). The grain yields were strongly supported by the yield contributing characters. It can be seen that yield increased upto a certain level of irrigation and then decreased. The results are in conformity with the findings of Talukder (1985), Chowdhury and Islam (1993) and Zirkov *et al.*, (1995). Different irrigation regimes were found to have significant effect on straw yields (Table 1). Straw yields

significantly increased and followed similar pattern as in grain yield. Chowdhury and Macksoud (1997) also found similar results.

The results on the yield and yield parameters of maize due to application of nitrogenous fertilizers are presented in Table 2. Significantly the highest plant height (269.40 cm) was found with 70 kg N/ha and was statistically identical (269.30 cm) with 100 kg N/ha. No significant differences were found between control and 120 kg N/ha treatments. Root length did not vary statistically due to application of nitrogenous fertilizers. Ear length and breath had marked variation due to changes in nitrogenous fertilizers (Table 2). Nitrogenous fertilizer significantly increased ears/plant while no variation was found in kernels/ear (Table 2). Highest number of ears/plant (1.33) was produced by 100 kg N/ha and no significant variation was found among 70 to 120 kg N/ha while the lowest (1.17) was in control. Different nitrogen treatments had significant influence on 100 kernel weight. The highest weight of 100 – kernel (28.49 g) was found with 100 kg N/ha and the lowest (27.89 g) with 120 kg N/ha. The grain yields were significantly influenced by different doses of nitrogen (Table 2). The grain yield was the highest

Table 2. Effect of nitrogen regimes on the yield and yield contributing characters of maize.

Nitrogen (kg N/ha)	Plant height (cm)	Root length (cm)	Ear length (cm)	Ear breath (cm)	Ears/plant (no.)	Kernels/Ear (no.)	100 Kernel weight (g)	Grain yield (t/ha)	Stover yield (t/ha)
Control	254.20b	28.75	18.78c	4.27c	1.17b	332.72	28.44ab	4.93b	8.21b
70	269.40a	28.38	18.88bc	4.51ab	1.31ab	341.78	27.90b	5.61a	8.88a
100	269.30a	28.91	19.21a	4.32bc	1.33a	344.37	28.49a	5.46a	8.89a
120	258.70b	28.91	19.12ab	4.69a	1.33a	336.34	27.89b	5.44a	8.99a
S _x	2.20	-	0.08	0.05	0.04	-	0.19	0.07	0.10
Level of significance	0.01	NS	0.01	0.01	0.01	NS	0.05	0.01	0.01

Note : Figure in a column having common letter(s) do not differ significantly but dissimilar letter differ significantly; NS = Not significant.

Table 3. Interaction effect of irrigation and nitrogen on the yield and yield contributing parameters of maize.

Interaction (Irrigation x Nitrogen)	Plant height (cm)	Root length (cm)	Ear length (cm)	Ear breath (cm)	Ears/plant (no.)	Kernel s/ear (no.)	100 Kernel weight (g)	Grain yield (t/ha)	Stover yield (t/ha)
I ₀ N ₀	240.80j	25.87	18.22efg	4.03ef	1.13	307.17	24.67ij	3.18k	5.42j
I ₀ N ₁	255.60ghij	25.30	18.42efg	3.85f	1.07	308.02	23.71ij	4.52ij	7.52ghi
I ₀ N ₂	252.7hij	26.80	18.76def	4.07ef	1.20	318.26	24.28ij	4.22j	7.14i
I ₀ N ₃	250.00ij	27.73	18.44efg	4.22cdef	1.07	307.85	24.93hij	3.50k	7.26hi
I ₁ N ₀	247.50ij	29.00	19.29bcd	4.72abc	1.13	397.43	25.56ghi	5.28efgh	8.75ef
I ₁ N ₁	269.50bcdefg	29.53	19.23cd	5.03ab	1.40	354.95	26.63fgh	5.80cdef	9.40de
I ₁ N ₂	273.60bcde	30.40	20.00ab	4.67abcd	1.40	365.94	26.63fgh	5.61def	9.69cd
I ₁ N ₃	273.00bcdef	29.07	20.37a	5.10a	1.47	371.70	27.96ef	6.41bc	10.27cd
I ₂ N ₀	258.30efghi	29.27	19.89abc	4.62abcd	1.27	345.55	31.15bc	6.05cd	10.49bc
I ₂ N ₁	274.80bcd	28.93	20.11a	4.93ab	1.47	367.25	31.05bc	7.26a	11.44a
I ₂ N ₂	279.70ab	28.67	20.54a	4.63abcd	1.40	350.97	34.00a	6.79ab	11.16ab
I ₂ N ₃	260.80defghi	28.60	20.31a	4.95ab	1.40	340.10	30.79bc	6.98ab	11.43a
I ₃ N ₀	268.90bcdefgh	30.67	18.68def	3.97ef	1.27	306.59	31.94b	5.07fghi	8.46fg
I ₃ N ₁	290.40a	29.47	18.30efg	4.13def	1.33	341.36	31.32bc	5.94cde	8.41fg
I ₃ N ₂	277.90abc	29.87	17.97fg	4.33cdef	1.47	350.39	31.00bc	5.95cde	8.72ef
I ₃ N ₃	257.00fghij	28.00	18.42efg	4.50bcde	1.47	334.22	29.94cd	5.47defg	7.82fghi
I ₄ N ₀	255.20ghij	28.93	17.80g	4.03ef	1.07	306.87	28.90de	5.08fghi	7.91fghi
I ₄ N ₁	256.40ghij	28.67	18.32efg	4.60abcd	1.27	337.34	26.80fg	4.55hij	7.62ghi
I ₄ N ₂	262.80cdefghi	28.80	18.81de	3.90f	1.20	336.31	26.57fgh	4.71hij	7.74ghi
I ₄ N ₃	252.70hij	31.13	18.07efg	4.67abcd	1.27	327.83	25.86ghi	4.86ghij	8.20fgh
S _x	4.91	-	0.18	0.12	-	-	0.42	0.17	0.22
Level of significance	0.05	NS	0.01	0.01	NS	NS	0.01	0.01	0.01

Note : Figure in a column having common letter(s) do not differ significantly but dissimilar letter differ significantly; NS = Not significant.

I₀ = Control, I₁ = IW/CPE ratio 0.2, I₂ = IW/CPE ratio 0.5, I₃ = IW/CPE ratio 0.8,

I₄ = IW/CPE ratio 1.0

N₀ = Control, N₁ = 70 kg N/ha, N₂ = 100 kg N/ha, N₃ = 120 kg N/ha.

(5.61 t/ha) due to the application of 70 kg N/ha and was statistically similar up to 120 kg N/ha. The lowest grain yield (4.93 t/ha) was produced by control. The results are in agreement with the findings of Cox *et al.*, (1993) and Gupta and Gautam (1994). Straw yield followed the similar pattern as in grain yield (Table 2). This might be due to the exuberant vegetative growth noted in the case of higher doses of nitrogen application.

Interaction effect of irrigation and nitrogen on the

performance of maize yield and yield contributing parameters are presented in Table 3. The highest grain yield (7.26 t/ha) was obtained by the combination of IW/CPE ratio of 0.5 and 70 kg N/ha while the lowest (3.81 t/ha) was in control. It is significantly supported by the yield contributing characters (Table 3). These results are in conformity with the findings of El-Noemami *et al.*, (1990) and Gab-Alla *et al.*, (1995). In combination, of irrigation and nitrogen had no significant effect on root length, ears/plant and

kernels/ear, respectively (Table 3).

The moisture status of the experimental plots during growing period of maize is shown in Fig. 1 to 5. Percentage of soil moisture gradually decreased through the soil profile in control (Fig. 1). Moisture status at 20 and 40cm depths of soil decreased days after irrigation and then gradually increased at every time of irrigation applied (Fig. 2). It might be due to elapsed time of percolation through the soil profile. In Fig. 3, 4 and 5 soil moisture status at 40 and 60cm depths of soil followed similar trend as in Fig. 2. But




Fig. 2. Soil moisture status at I₁ (IW/CPE=0.2).




Fig. 1. Soil moisture status at I₀ (IW/CPE=0.0).




Fig. 3. Soil moisture status at I₂ (IW/CPE=0.5).



Fig. 4. Soil moisture status at I₃ (IW/CPE=0.8).Fig. 5. Soil moisture status at I₄ (IW/CPE=1.0).

Table 4. Physico-chemical properties of soil during land preparation.

Treatment	pH	EC (µs/cm)	N (%)	NH ₄ -N (ppm)	NO ₃ -N (ppm)	P (ppm)	K (ppm)	S (ppm)	Organic carbon (%)	Soil texture
I ₀ N ₀	6.48	420	0.162	4.20	11.20	57	42	20	0.95	Silt loam
I ₀ N ₁	6.43	406	0.128	8.40	14.00	59	44	15	1.00	"
I ₀ N ₂	7.42	440	0.055	7.00	12.60	57	40	15	1.00	"
I ₀ N ₃	6.60	464	0.026	7.00	9.80	59	40	34	0.48	"
I ₁ N ₀	7.72	478	0.124	5.60	4.20	50	50	20	1.48	"
I ₁ N ₁	6.76	430	0.059	3.80	7.00	58	64	25	0.84	"
I ₁ N ₂	6.78	340	0.028	4.20	4.80	56	62	15	1.48	"
I ₁ N ₃	7.80	368	0.008	5.80	4.80	59	66	20	1.26	"
I ₂ N ₀	6.74	488	0.052	4.20	7.00	60	48	20	0.79	"
I ₂ N ₁	6.86	354	0.124	4.20	4.20	69	40	28	1.44	"
I ₂ N ₂	6.87	468	0.139	4.20	8.40	59	44	28	0.61	"
I ₂ N ₃	7.60	401	0.152	3.80	8.40	64	40	18	0.79	"
I ₃ N ₀	6.67	440	0.131	5.60	8.40	56	39	20	0.53	"
I ₃ N ₁	7.48	356	0.047	5.60	12.60	59	62	30	0.70	"
I ₃ N ₂	6.76	483	0.039	4.20	8.40	60	39	30	1.61	"
I ₃ N ₃	7.23	340	0.068	8.00	8.67	58	50	28	0.88	"
I ₄ N ₀	7.80	402	0.990	7.68	10.11	50	55	28	0.79	"
I ₄ N ₁	7.68	411	0.101	8.10	6.67	60	60	30	0.90	"
I ₄ N ₂	7.70	414.35	0.800	9.20	10.45	62	48	35	0.69	"
I ₄ N ₃	7.52	378	0.850	8.88	8.85	55	56	27	1.00	"
Mean	7.14	414.35	0.168	5.98	8.68	58.35	49.45	24.30	0.96	
Range	6.42-7.80	340-488	0.068-0.162	3.80-8.88	4.80-14.00	50-69	39-66	15-34	0.48-1.61	

Table 5. Physico-chemical properties of soil at harvest.

Treat- ment	pH	EC ($\mu\text{s}/\text{cm}$)	N (%)	$\text{NH}_4\text{-N}$ (ppm)	$\text{NO}_3\text{-N}$ (ppm)	P (ppm)	K (ppm)	S (ppm)	Organic carbon (%)
I_0N_0	6.52	411	0.131	8.40	10.00	49	27	17	0.88
I_0N_1	6.88	434	0.096	8.40	12.60	55	28	13	1.02
I_0N_2	6.90	419	0.086	8.40	19.60	58	29	14	0.93
I_0N_3	7.03	439	0.075	9.80	4.20	53	60	32	0.83
I_1N_0	7.12	448	0.077	7.00	8.40	52	67	13	0.85
I_1N_1	7.04	402	0.085	11.20	4.80	55	68	14	0.95
I_1N_2	7.16	376	0.087	8.40	7.00	51	47	19	0.95
I_1N_3	6.99	407	0.064	7.00	8.40	54	39	15	0.74
I_2N_0	6.73	337	0.121	5.60	4.80	56	36	16	1.14
I_2N_1	7.12	381	0.141	7.00	5.60	59	43	30	1.28
I_2N_2	6.57	382	0.064	7.00	19.00	55	39	27	0.72
I_2N_3	6.58	479	0.072	7.00	14.00	59	39	17	0.79
I_3N_0	6.98	394	0.062	7.00	7.00	56	40	15	0.67
I_3N_1	6.61	327	0.078	5.60	11.20	50	48	24	0.85
I_3N_2	6.89	470	0.082	11.20	5.60	49	64	22	0.93
I_3N_3	7.30	312	0.088	8.30	10.50	38	30	22	0.78
I_4N_0	7.22	372	0.120	8.88	11.60	42	37	28	0.89
I_4N_1	7.60	410	0.131	7.60	8.67	49	48	27	0.67
I_4N_2	7.13	375	0.088	7.89	7.89	37	51	33	0.88
I_4N_3	7.33	340	0.990	7.00	6.78	51	55	22	0.85
Mean	6.99	395.75	0.137	7.93	9.38	51.40	44.75	13-32	0.88
Range	6.52- 7.60	312- 479	0.064- 0.141	5.60- 11.20	4.20- 19.60	37-59	27-74	9-40	0.67-1.28

moisture status at 20 cm depth of soil fluctuated at prior to and post irrigation, respectively. It might be due to low infiltration, evaporation and water holding capacity of soil.

The physico-chemical properties of soil during land preparation are presented in Table 4. Texturally, the soil of the study area was silt loam. pH, electrical conductivity (EC), nitrogen (N), $\text{NH}_4\text{-N}$, $\text{NH}_3\text{-N}$, phosphorous (P), potassium (K), sulphur (S) and organic carbon varied from 6.42 to 7.80, 340 to 488 $\mu\text{s}/\text{cm}$, 0.068 to 0.162%, 3.80 to 8.88 ppm, 4.80 to 14.00 ppm, 50 to 69 ppm, 39 to 66 ppm, 15 to 34 ppm and 0.48 to 1.16%, respectively. The physico-chemical properties of soil at harvest are presented in Table 5. It was found that pH, electrical conductivity (EC), nitrogen (N), $\text{NH}_4\text{-N}$, $\text{NH}_3\text{-N}$, phosphorous (P), potassium (K), sulphur (S) and organic carbon varied

from 6.52 to 7.60, 312 to 479 $\mu\text{s}/\text{cm}$, 0.064 to 0.141%, 5.60 to 11.20 ppm, 4.20 to 19.60 ppm, 37 to 59 ppm, 27 to 74 ppm, 13 to 32 ppm and 0.67 to 1.28%, respectively. Table 4 and 5 indicate that the plant nutrients N, P, K, S and percentage of organic carbon decreased slightly at harvest in comparison with land preparation. This might be due to the uptake of plant nutrients during the growing period of the crop.

ABSTRACT

An experiment was carried out during November 1998 to April 1999 to find out the response of yield and yield contributing parameters of maize (cv. Bornali) to water stress and nitrogenous fertilizer. The experiment included two factors such as five irrigation regimes and four nitrogen levels. Texturally, the soil was silty loam. Yield and yield contributing characters were

significantly affected due to the application of irrigation and nitrogen. The highest grain yield of 6.77 t/ha was obtained with IW/CPE ratio of 0.5 and 5.61 t/ha by the application of 70 kg N/ha. Interactions between IW/CPE ratio of 0.5 and 70 kg N/ha were the best combination for yield of maize.

水ストレスと窒素肥料がトウモロコシの生育と収量に及ぼす影響

要 旨

トウモロコシの水分ストレスと窒素肥料に対する反応を調べるために、1998年11月から1999年4月にかけて実験を行なった。実験は、灌漑水量5水準と4水準の窒素レベルについて行なった。実験にはシルト質ロームを用いた。灌漑水量と窒素レベルにより収量は明らかな影響を受けた。灌漑水量に関して、最大収量の6.77t/haはIW/CPEが0.5で得られた。また、窒素レベルに関しては、最大収量の5.61t/haは70kgN/haの窒素施肥で得られた。従って、IW/CPE=0.5の灌漑水量と70kgN/haの窒素施肥量は、トウモロコシ栽培において高収量を得るのに良い組み合わせであるといえる。

REFERENCES

- Abrecht, D. G. and P. S. Carberry 1993. The influence of water deficit prior to tassel initiation on maize growth, development and yield. *Field Crop Res.* 31(1-2): 55-69.
- BARC 1997. Fertilizer Recommendation Guide. Bangladesh Agricultural Research Council, Farm Gate, Dhaka. p 45.
- BBS 1997. Year book of Agricultural Statistics of Bangladesh, Bureau of Statistics Division, Ministry of Planning, Dhaka, Bangladesh. p.164.
- Chowdhury, M. K. and M. A. Islam 1993. Production and use of maize(in Bengali). Edited by Chowdhury and Islam. On-Farm Research Division. Bangladesh Agricultural Research institute, Joydebpur, Gazipur. pp. 8 &57.
- Chowdhury, N. H. and S. W. Macksoud 1997. Effect of irrigation schedule and plant population on the grain yield and other plant characters of hybrid corn grown in Bequa's Plain, Lebanon. *West Pak J. Agril. Res.* 5(2): 77-79.
- Cox, W. J., S. Kalonge, D. J. R. Cherney and W. S. Raid 1993. Growth, yield and quality of forage maize under different nitrogen management practices. *Agron. J.* 85(2): 341-347.
- Dai, J. Y., W. L. Gu, X. Y. Shen, B. Zheng, H. Qi and S. F. Cai 1990. Effect of drought on the development and yield of maize at different growth stages. *J. Shenyang Agril. Univ.* 21(3): 181-185.
- El-Noemani, A. A., A. K. A. EI-Halim, H. A. EI-Zeiny and A. K. Abd-EI-Halim 1990. Response of maize to irrigation intervals under different levels of nitrogen fertilization. *Egypt J. Agron.* 15(1-2): 147-158.
- Gab-Alla, F. I., M. M. Shahin, H. M. Eid and S. M. EL-Marsafawy 1995. Relationship between nitrogen level, yield and its components through some irrigation regimes for maize. *Egypt J. Soil Sc.* 35(3): 297-310.
- Gomez, K. A. and A. A. Gomez 1984. Statistical procedure for agricultural research. Second Edn. International Rice Research Institute. Los Banos, Manila, The Philippines. pp. 204-207.
- Gordon, W. B., R. J. Raney and L. R. Stone 1995. Irrigation management practice for corn

- production in north central Kansas. *J. Soil and Water Cons.* 50(4): 395 - 398.
- Gupta, D. K. and R. C. Gautam 1994. Response of maize to nitrogen rate and schedule of tasseling removal. *Indian J. Ecol.* 21(1): 73 - 74.
- Ne Smith, D. S. and J. T. Ritche 1992. Short and long-term responses of corn to a pre-anthesis soil water deficit. *Agron. J.* 84(1): 107 - 113.
- Singh, K. K., A. Fazeel and R. C. Gautam 1996. Response of maize to nitrogen and percentage of tassel removal. *Annals Agric. Res.* 17(2): 213 - 214.
- Shoazhong, K. and Z. Mingannang 1992. Crop water production function of maize for Northeast Brazil. *Pesquisa Agropecuaria Brasileira.* 23(12): 1413 - 1420.
- Talukder, M. S. U. 1985. Yield response of sweet corn to irrigation and fertilization grown in north-eastern Thailand. *Bangladesh J. Agril. Sc.* 10(2): 87 - 95.
- Zhirkov, Z. V. 1995. Growing maize for grain under optimum and deficit irrigation. *Rasteniev "dni-Nauki".* 32(9 - 10): 142 - 145.