

琉球大学学術リポジトリ

Surgical outcomes of displaced proximal humeral fractures : antegrade intramedullary nail versus locking plate

メタデータ	言語: 出版者: 琉球医学会 公開日: 2020-11-24 キーワード (Ja): キーワード (En): proximal humeral fractures, antegrade intramedullary nails, locking plates, avascular necrosis, varus deformity 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/20.500.12000/0002016913

Surgical outcomes of displaced proximal humeral fractures: antegrade intramedullary nail versus locking plate

Isoya Goya^{1, 2)}, Hiroshi Yamaguchi^{2, 3)}, Takashi Toma²⁾,
Tomohiro Moriyama⁴⁾, Fuminori Kanaya²⁾

¹⁾ *Department of Orthopedic Surgery, Yonabaru Chuo Hospital*

²⁾ *Department of Orthopedic Surgery, Graduate School of Medicine, University of the Ryukyus*

³⁾ *Rehabilitation Clinic Yamaguchi*

⁴⁾ *Okinawa Red Cross Hospital*

(Received on January 15, 2019, accepted on February 4, 2019)

ABSTRACT

This study aimed to compare the postoperative shoulder range of motion (ROM) and complication rates in patients with proximal humeral fractures that were treated with antegrade intramedullary nails or locking plates. Between 2008 and 2016, the shoulders of 108 patients with proximal humeral fractures underwent internal fixation using antegrade intramedullary nails (N group) or locking plates (P group). Of these, 42, 48, and 18 shoulders exhibited two-, three-, and four-part fractures, respectively. For patients with three- or four-part fractures, the mean age in the P group was significantly less compared with that in the N group. The mean operation time, blood loss, and postoperative follow-up period did not significantly differ between the N and P groups. The external rotation of two-part fractures in the N group was significantly better than in the P group. The external rotation of three- and four-part fractures in patients aged 65–74 years was significantly better in the N group than in the P group. For two-part fractures in the N group, forward flexion was better in patients aged <65 years than in those aged ≥65 years. In three- and four-part fractures, forward flexion was better in patients aged 65–74 years than in those aged ≥75 years in the N group, and it was better in patients aged <65 years than in those aged ≥75 years in the P group. In three- and four-part fractures, external rotation was better in patients aged 65–74 years than in those aged ≥75 years in the N group and better in patients aged <65 years than in those aged ≥65 years in the P group. Only one patient with a three-part fracture in the P group exhibited nonunion. Avascular necrosis was more likely to occur in patients with three- and four-part fractures. Varus deformity was likely to frequently occur in patients with two-part fractures in the P group. Surgical treatment of proximal humeral fractures provided comparable clinical results in the N and P groups. Except for external rotation, patients with two-part fractures in the N group showed better outcomes than those in the P group. Moreover, elderly patients exhibited poor ROM. *Ryukyu Med. J., 38 (1~4) 61~72, 2019*

Key words: proximal humeral fractures, antegrade intramedullary nails, locking plates, avascular necrosis, varus deformity

INTRODUCTION

Proximal humeral fractures comprise the second most common type of fracture of the upper extremities in the elderly and account for 5% of all

fractures in adults^{1, 2)}. Particularly, adults aged ≥60 years are vulnerable to complex three- and four-part humeral fractures³⁾. In 2005, approximately 12.8 million Japanese individuals suffered from osteoporosis and this number is gradually increasing⁴⁾. Most patients with non-displaced or

Corresponding Author: Isoya Goya, Department of Orthopedic Surgery, Graduate School of Medicine, University of the Ryukyus, 207 Uehara, Nishihara 903-0215, Okinawa, Japan. Tel: +81-98-895-1174; Fax: +81-98-895-1424, E-mail: isoppe581005@yahoo.co.jp

minimally displaced fractures are conservatively treated. However, displaced proximal humeral fractures often require surgical treatment, for which several surgical instruments have been developed.

Antegrade intramedullary nails or locking plates are the most commonly used surgical instruments for treating such fractures. However, achieving stable fixation is often difficult due to the comminuted medial fragment and osteoporosis background, particularly in the elderly. Several previous studies have assessed the use of antegrade intramedullary nails and locking plates for treating proximal humeral fractures⁵⁻¹¹. However, there is no clear consensus regarding the optimal management of proximal humeral fractures.

The present study aimed to compare the postoperative shoulder range of motion (ROM) and complication rate in patients with proximal humeral fractures that were treated using antegrade intramedullary nails and locking plates at the Ryukyus University Hospital and affiliated hospitals. Fourteen experienced orthopedic surgeons were involved in this study, and the patients in this study are most likely to represent patients with proximal humeral fractures who underwent surgical treatment in Okinawa.

PATIENTS AND METHODS

Patient selection

Between 2008 and 2016, 108 proximal humeral fractures in 108 patients were treated with internal fixation using antegrade intramedullary nails or locking plates. The exclusion criteria included patients with open fractures, neurological injury-

related fractures, and pathological fractures. All patients agreed to participate were included in the study, postoperative follow-up period were at least 12 months. The research protocol was approved by the Institutional Review Board of our institute (No. 388), and all patients provided written informed consent prior to initiating the study.

All procedures were conducted at the Ryukyus University Hospital and affiliated hospitals by one of 14 experienced orthopedic surgeons. The use of antegrade intramedullary nails or locking plates was dependent on the surgeon's preference. The 108 shoulders were categorized into two groups based on the method of internal fixation: N (antegrade intramedullary nails) and P (locking plates) groups (54 patients per group). Anteroposterior radiographs were used to assess the fracture type using the Neer classification. Accordingly, 42, 48, and 18 shoulders were found to have two-, three-, and four-part fractures, respectively. Cases were grouped as two-part fractures or three- and four-part fractures.

N group

Antegrade intramedullary nail fixation was performed using the New Straight Nail System[®] (Teijin Nakashima Medical, Okayama, Japan). The N group comprised 11 men and 43 women. Two-part fractures were found in 23 shoulders (5 men and 18 women) and three- and four-part fractures in 31 shoulders (6 men and 25 women). The mean patient age at operation was 67 (range, 35–91) years for two-part fractures and 70 (range, 46–85) years for three- and four-part fractures. The mean follow-up period were 18 (range, 12–78) and 24 (range, 12–64) months for two-part and three- and four-part fractures, respectively (Table 1).

Table 1 Demographic characteristics of the study population

	Two-part fractures (n=42)			Three and four-part fractures (n=66)		
	Antegrade intramedullary nails	Locking plates	P value	Antegrade intramedullary nails	Locking plates	P value
Number of patients (men/women)	23 (5/18)	19 (3/16)	0.93	31 (6/25)	35 (12/23)	0.28
Average age (years, range)	67 (35–91)	66 (21–87)	0.62	70 (46–85)	61 (20–83)	<0.01
Age, <65 years	10 (44%)	6 (32%)		6 (18%)	23 (64%)	
Age, 65–74 years	7 (30%)	4 (21%)		16 (52%)	6 (21%)	
Age, ≥ 75 years	6 (26%)	9 (47%)		9 (33%)	6 (15%)	
Follow-up (months, range)	18 (12–78)	19 (12–48)	0.24	24 (12–64)	23 (12–114)	0.53

* P<0.05

P group

The locking plate fixation was performed using the PHILOS® System (DePuy-Synthes, Solothurn, Switzerland). The P group comprised 15 men and 39 women. Two-part fractures were found in 19 shoulders (3 men and 16 women) and three- and four-part fractures in 35 shoulders (12 men and 23 women). The mean patient age at operation was 66 (range, 21–87) years for two-part fractures and 61 (range, 20–83) years for three- and four-part fractures. The mean follow-up period were 19 (range, 12–48) and 23 (range, 12–114) months for two-part fractures and three- and four-part fractures, respectively (Table 1).

Surgical technique

In both groups, surgery was performed under general anesthesia with an interscalene nerve block with the patient placed in the beach-chair position. After reduction, antegrade intramedullary nail or locking plate was fixed. For patients with large bone defects, the iliac bone or alpha-tricalcium phosphate paste (artificial bone) with local cancellous bone chips was grafted.

N group

A longitudinal incision was made using the through anterolateral transdeltoid approach in the supraspinatus muscle by splitting the anterior and

middle thirds of the deltoid muscle. Comminuted fractures were reduced using the deltopectoral approach. The nail was inserted near the top of the humeral head, which inevitably damages the cartilage of the humeral head. After nail insertion, a proximal interlocking screw was inserted from the lateral aspect of the humeral head. Subsequently, two or three distal interlocking screws were inserted at the deltoid insertion to avoid radial nerve injuries. Another proximal interlocking screw (from the anterior to posterior direction) and an end cap were also placed. Finally, the incised rotator cuff was closed using non-absorbable sutures (Figure 1).

P group

A longitudinal incision was made using the deltopectoral approach. Locking plates were placed in a lateral position to the bicipital groove and 1 cm inferior to the upper humeral greater tuberosity, laterally and posteriorly. Guidewires were inserted through the proximal holes of the plate, followed by distal fixation using cortical or locking screws. At least five locking screws were inserted into the humeral head; moreover, two locking screws were inserted to the humerus diaphysis (Figure 2).

Rehabilitation protocol

The postoperative rehabilitation protocol was similar for both N and P groups. Postoperatively,



(a) preoperative radiograph



(b) preoperative 3D computed tomography



(c) postoperative radiograph (13 months after operation)

Fig.1 70-year-old female who fell over and was suffered from a two-part proximal humerus fracture. Antegrade intramedullary nail fixation was performed.



(a) preoperative radiograph



(b) preoperative 3D computed tomography



(c) postoperative radiograph (16 months after operation)

Fig.2 55-year-old male who fell from height and was suffered from a four-part proximal humerus fracture. Locking plate fixation was performed.

patient arms were immobilized using a sling and/or abduction pillow for 4–6 weeks. Postoperative passive ROM exercises were initiated 1–3 weeks postoperatively, and active ROM exercises were initiated 4–6 weeks postoperatively. For patients with three- or four-part fractures, longer durations of immobilization and slower ROM exercises were adopted.

Radiographic assessment

Shoulders were radiographically evaluated in the following three views: anteroposterior view in the scapular plane, lateral view of the scapula, and supine axillary view. Radiographs were obtained immediately after operation and at follow-up.

Radiographs were evaluated for bone union and complications. Any complications observed during the entire follow-up period were recorded, including fracture nonunion, avascular necrosis, varus deformity, and screw back-out. Union was defined as visible continuity of the cortical bone that could be assessed using at least two radiographic views. Postoperative avascular necrosis was classified using the Cruess classification¹²⁾. Varus deformity was considered in cases with $>20^\circ$ deformity.

Clinical assessment

Active ROM was evaluated using forward flexion and external rotation, operation time, and blood loss in the N and P groups and then compared

between subgroups classified by the type of fracture (two-part versus three- or four-part fractures). Patients were also classified into three age groups (<65 , $65-74$, or ≥ 75 years).

Statistical analysis

Descriptive statistics of the clinical data were determined. The Kruskal–Wallis test and Mann–Whitney test were used to determine statistical significance for quantitative variables, whereas the Fisher exact test was used for qualitative variables. A P-value of <0.05 was considered to be statistically significant.

RESULTS

For patients with three- and four-part fractures, the mean age in the P group was significantly less compared with that in the N group ($P < 0.01$). Gender and mean postoperative follow-up period did not significantly differ between the two groups (Table 1). Furthermore, operation time, blood loss, and mean postoperative follow-up period were not significantly different between the groups. The external rotation of two-part fractures in the N group was significantly better than that of those in the P group ($P < 0.02$) (Table 2). Moreover, for patients aged 65–74 years, the external rotation of three- and four-part fractures in the N group was significantly

Table 2 Active range of motion in forward flexion and external rotation in the N and P groups disaggregated by the type of fracture

	Two-part fractures			Three and four-part fractures		
	Antegrade intramedullary nails, Mean \pm SD	Locking plates, Mean \pm SD	P value	Antegrade intramedullary nails, Mean \pm SD	Locking plates, Mean \pm SD	P value
Forward flexion ($^{\circ}$)	127 \pm 26.82	114 \pm 29.06	0.16	117 \pm 25.11	116 \pm 36.67	0.69
External rotation ($^{\circ}$)	39 \pm 17.06	27 \pm 11.93	<0.02	35 \pm 22.70	31 \pm 16.49	0.36
Operation time (minutes)	146 \pm 77.68	148 \pm 27.17	0.54	176 \pm 79.45	143 \pm 32.22	0.08
Blood loss (ml)	172 \pm 200.56	203 \pm 83.52	0.42	198 \pm 158.27	249 \pm 154.52	0.36

* P<0.05

The external rotation of two-part fractures in the antegrade intramedullary nails group was significantly better than that of those in the locking plates group

Table 3 Active range of motion in forward flexion and external rotation post operation in the antegrade intramedullary nails and locking plates disaggregated by the type of fracture and various age subgroups

		Two-part fractures			Three and four-part fractures		
		Antegrade intramedullary nails, Mean \pm SD	Locking plates, Mean \pm SD	P Value	Antegrade intramedullary nails, Mean \pm SD	Locking plates, Mean \pm SD	P Value
Forward flexion in patients aged: ($^{\circ}$)	<65 years	144 \pm 19.85	128 \pm 30.91	0.34	119 \pm 19.02	123 \pm 34.00	0.72
	65–74 years	120 \pm 14.14	119 \pm 15.16	0.91	127 \pm 23.57	115 \pm 30.41	0.46
	\geq 75 years	108 \pm 30.79	103 27.99	0.81	100 \pm 22.11	83 \pm 36.70	0.39
External rotation in patients aged: ($^{\circ}$)	<65 years	42 \pm 18.48	31 \pm 10.57	0.21	39 \pm 17.42	36 \pm 16.70	0.75
	65–74 years	38 \pm 9.95	25 \pm 11.18	0.15	42 \pm 18.71	15 \pm 10.00	<0.001
	\geq 75 years	38 \pm 19.09	24 \pm 9.13	0.21	20 \pm 24.94	24 \pm 7.31	0.67

* P<0.05


For patients aged 65–74 years, the external rotation of three- and four-part fractures in the antegrade intramedullary nails group was significantly better than that of those in the locking plates group.

better than that of those in the P group (P<0.001). Forward flexion was not significantly different between the N and P groups (Table 3).

Among patients with two-part fractures in the N group, forward flexion was better in those aged <65 years than in those aged \geq 65 years (Figure 3A). Among patients with three- and four-part fractures in the N group, forward flexion was better in patients aged 65–74 years than in patients aged \geq 75 years. Among patients with three- and four-part fractures in the P group, forward flexion was better in patients aged <65 years than in patients aged \geq 75 years (Figure 3B). External rotation among


patients with two-part fractures was comparable in both groups (Figure 4A). Among patients with three- and four-part fractures in the N group, external rotation was better in patients aged 65–74 years than in patients aged \geq 75 years. Among patients with three- and four-part fractures in the P group, external rotation was better in patients aged <65 years than in patients aged \geq 65 years (Figure 4B).

Only one patient with a three-part fracture in the P group exhibited nonunion. Avascular necrosis occurred more frequently in patients with three- and four-part fractures (13 patients, 20%) than in those with two-part fractures (1 patient, 2%; P<0.02). Avascular necrosis occurred in six (60%)




In the N group, forward flexion was better in those aged <65 years than in those aged ≥65 years.

Fig.3A Active postoperative range of motion in forward flexion in patients with two-part fractures in the antegrade intramedullary nail group (N group) and locking plate group (P group) disaggregated by various age groups




In the N group, forward flexion was better in patients aged 65-74 years than in patients aged ≥75 year. In the P group, forward flexion was better in patients aged <65 years than in patients aged ≥75 years.

Fig.3B Active postoperative range of motion in forward flexion in patients with three- and four-part fractures in the nail group (N group) and locking plate group (P group) disaggregated by various age groups



No significant differences were observed in the N and P groups among patients of different age groups.

Fig. 4A Active postoperative range of motion in external rotation in patients with two-part fractures in the nail group (N group) and locking plate group (P group) by age groups



In the N group, external rotation was better in patients aged 65-74 years than in patients aged ≥75 years. In the P group, external rotation was better in patients aged <65 years than in patients aged ≥65 years.

Fig. 4B Active postoperative range of motion in external rotation in patients with three- and four-part fractures in the nail group (N group) and locking plate group (P group) disaggregated by age groups

and four (50%) patients with four-part fractures in both the N and P groups, respectively. Varus deformity frequently occurred in patients with two-part fractures in the P group (four patients, 21%). Screw back-out was observed in one patient in the N group with a two- and three-part fracture each and in one patient in the P group with a four-part fracture. Re-displacement of the reduced humeral head was performed in one patient with a four-part fracture in the N and P group each; both patients experienced severe pain due to avascular necrosis (Table 4).

DISCUSSION

Stable fixation of displaced proximal humeral fractures in the elderly is often difficult owing to the presence of osteoporosis. Severe comminution and weak bone, which inhibit rigid fixation, are common manifestations of osteoporotic proximal humeral fractures. Recently, several instruments have been developed for ensuring rigid fixation of proximal humeral fractures. Among these, antegrade intramedullary nails and locking plates are considered as the primary instruments^{9, 10, 13-18}. Studies from Western countries have demonstrated comparable clinical outcomes with the use of antegrade intramedullary nails and locking plates^{10, 13-18}. However, no studies have compared the outcomes using the aforementioned instruments in elderly Japanese individuals with proximal humeral fractures. Thus, the present study investigated

surgically treated proximal humeral fractures in elderly patients at the Ryukyus University Hospital and affiliated hospitals and compared the clinical outcomes associated with the use of antegrade intramedullary nails and locking plates.

Compared with locking plates, antegrade intramedullary nails are supposed to have the advantages of being less invasive and causing less damage to the blood supply of the humeral head¹⁰. In addition, antegrade intramedullary nails ensure more rigid fixation than locking plates. However, antegrade intramedullary nails have the disadvantages of incising the rotator cuff and damaging the cartilage of the humeral head when inserting nails^{13, 14, 19}. Reportedly, locking plates have the advantages of greater anatomical reduction and rigid fixation of comminuted fragments. However, their disadvantages include extensive soft tissue dissection, longer operation time, more blood loss, and mechanical impingement with the acromion^{20, 21} (Table 5).

In the present study, the degree of external rotation that was achieved in the P group was significantly lower than that achieved in the N group for three- or four-part fractures. Locking plates are typically implanted via the deltopectoral approach by incising the anterior surface of the shoulder. The deltopectoral approach is considered to be more invasive than the anterolateral approach for inserting antegrade intramedullary nails. The invasive approach can cause tissue coalescence, resulting in restrained external rotation. During the postoperative period, patients treated using locking

Table 4 Postoperative complications

	Antegrade intramedullary nails			Locking plates		
	Two-part fractures (N=23)	Three-part fractures (N=21)	Four-part fractures (N=10)	Two-part fractures (N=19)	Three-part fractures (N=27)	Four-part fractures (N=8)
Non-union	0	0	0	0	1 (4%)	0
Avascular necrosis	1 (4%)	2 (10%)	6 (60%)	0	1 (4%)	4 (50%)
Varus deformity	1 (4%)	1 (5%)	0	4 (21%)	2 (7%)	1 (13%)
Screw back-out	1 (4%)	1 (5%)	0	0	0	1 (13%)
Re-displacement	0	0	1 (10%)	0	0	1 (13%)

One patient with a three-part fracture in the locking plates developed nonunion. Avascular necrosis was more likely to occur in patients with four-part fractures. Varus deformity frequently occurred in patients with two-part fractures in the locking plates group.

Table 5 Advantages and disadvantages of the antegrade intramedullary nails and locking plates

	Antegrade intramedullary nail	Locking plate
Advantages	<ul style="list-style-type: none"> • less invasion • rigid fixation 	<ul style="list-style-type: none"> • more anatomical reduction
Disadvantages	<ul style="list-style-type: none"> • rotator cuff incision • cartilage damage of humeral head 	<ul style="list-style-type: none"> • more invasion • more operation time and blood loss • mechanical impingement

plates experienced two- to three-fold greater pain than those treated with antegrade intramedullary nails¹⁸); this may be due to the more invasive nature of the surgery required for treatment using locking plates.

A meta-analysis by Li *et al.* demonstrated that operation time and perioperative blood loss associated with the use of antegrade intramedullary nails were less compared with those associated with the use of locking plates for two-, three-, and four-part fractures²². In the present study, the operation time and perioperative blood loss did not differ between the N and P groups. This could be because the experienced surgeons in this study preferred using locking plates.

Our results also showed that elderly patients had poorer ROM than younger patients. This is consistent with other reports demonstrating that elderly patients tend to exhibit poorer outcomes due to osteoporosis^{10, 15, 16}.

Avascular necrosis and varus deformity are two major complications associated with the surgical treatment of proximal humeral fractures. Avascular necrosis occurs due to humeral head ischemia caused by fractures, leading to screw extrusion at the humeral head. In our study, avascular necrosis occurred in 22.6% and 14.3% of patients with three- and four-part fractures in the N and P groups, respectively; this difference was not statistically significant. Avascular necrosis occurred more frequently in patients with three- or four-part fractures (13 patients, 20%) than in two-part fractures (one patient, 2%). Boudard *et al.* reported that avascular necrosis occurred in 26.7% of antegrade intramedullary nails cases and 21.2% of locking plate cases for three- or four-part fractures¹⁷. Hertel *et al.* reported that the length of the dorsomedial metaphyseal extension, the integrity of the medial hinge, and the type of basic fracture determined using the binary description system are the best predictors of ischemia²³. In the present study, all 14 patients with avascular necrosis exhibited a high-risk fracture type

as defined by Hertel. Specifically, 10 patients showed a medial neck fragment shorter than 8 mm, and 4 patients exhibited disrupted medial hinge. These results suggest that humeral head replacement should be considered in high-risk cases.

Varus deformity in our study occurred most frequently in patients with two-part fractures in the P group. This is consistent with a report by Zhu *et al.* that demonstrated high rates of varus deformity among patients treated with locking plates for two-part fractures¹⁰. According to DePalma and Nobuhara, the permissible angle of varus-valgus deformation is up to 20°^{24, 25} and deformation greater than this angle can cause cuff dysfunction²⁶. A biomechanical study revealed that the use of antegrade intramedullary nails entailed greater stiffness for varus, valgus, and torsional loading than locking plates²⁷. Gardner *et al.* reported that placing a superiorly directed oblique locked screw in the inferomedial region of the proximal fragment helped achieve more stable medial column support in the locking plate²⁸. However, in our study, the oblique locked screw was inserted in 17 of 19 two-part fractures and did not seem to prevent varus deformities.

Reportedly, complications can be prevented to some extent by improving surgical techniques; for example, varus deformity decreases with the use of additional non-absorbable sutures tagged through the rotator cuff tendons or tuberosity fragments and fixed at the base of the blade, passed through holes in the plate, or fixed using additional screws^{29, 30}. These complications can also be prevented by maintaining an appropriate external fixation period and adopting a rehabilitation program based on the patient's age and fracture type³⁰.

LIMITATIONS

A higher number of cases is required to further evaluate the surgical results of fracture subgroups.

Our study design was neither randomized nor prospective. Further, the preferences of the surgeons to use intramedullary nails or locking plates may have affected the results.

CONCLUSIONS

In summary, the surgical treatment of proximal humeral fractures using antegrade intramedullary nails (N group) or locking plates (P group) produced comparable clinical results. Except for external rotation, patients with two-part fractures in the N group showed better outcomes than those in the P group. Elderly patients exhibited especially poor ROM. Avascular necrosis was more likely in patients with three- and four-part fractures.

DISCLAIMER

The authors, their immediate families, and any research foundation to which they are affiliated did not receive any funding or other benefits from any commercial entity related to the subject of this study.

REFERENCES

- 1) Palvanen M., Kannus P., Niemi S., Parkkari J.: Update in the epidemiology of proximal humeral fractures. *Clin Orthop Relat Res.* 442: 87-92, 2006.
- 2) Hak DJ., Mauffrey C., Hake M., Hammerberg EM., Stahel PF.: Ipsilateral femoral neck and shaft fractures: current diagnostic and treatment strategies. *Orthopedics.* 38(4): 247-251, 2015.
- 3) Bahrs C., Stojicevic T., Blumenstock G., Brorson S., Badke A., Stöckle U., Rolauuffs B., Freude T.: Trends in epidemiology and patho-anatomical pattern of proximal humeral fractures. *Int Orthop.* 38(8): 1697-1704, 2014.
- 4) Yoshimura N., Muraki S., Oka H., Kawaguchi H., Nakamura K., Akune T.: Cohort profile: research on Osteoarthritis/Osteoporosis Against Disability study. *Int J Epidemiol.* 39(4): 988-995, 2010.
- 5) Brunner F., Sommer C., Bahrs C., Heuwinkel R.,

- Hafner C., Rillmann P., Kohut G., Ekelund A., Muller M., Audigé L., Babst R.: Open reduction and internal fixation of proximal humerus fractures using a proximal humeral locked plate: a prospective multicenter analysis. *J Orthop Trauma.* 23(3): 163-172, 2009.
- 6) Thanasas C., Kontakis G., Angoules A., Limb D., Giannoudis P.: Treatment of proximal humerus fractures with locking plates: a systematic review. *J Shoulder Elbow Surg.* 18(6): 837-844, 2009.
- 7) Brorson S., Rasmussen JV., Frich LH., Olsen BS., Hróbjartsson A.: Benefits and harms of locking plate osteosynthesis in intraarticular (OTA Type C) fractures of the proximal humerus: a systematic review. *Injury.* 43: 999-1005, 2012.
- 8) Sproul RC., Iyengar JJ., Devcic Z., Feeley BT.: A systematic review of locking plate fixation of proximal humerus fractures. *Injury.* 42: 408-413, 2011.
- 9) Gradl G., Dietze A., Käab M., Hopfenmüller W., Mittlmeier T.: Is locking nailing of humeral head fractures superior to locking plate fixation? *Clin Orthop Relat Res.* 467 (11): 2986-2993, 2009.
- 10) Zhu Y., Lu Y., Shen J., Zhang J., Jiang C.: Locking intramedullary nails and locking plates in the treatment of two-part proximal humeral surgical neck fractures: a prospective randomized trial with a minimum of three years of follow-up. *J Bone Joint Surg Am.* 93: 159-168, 2011.
- 11) Hardeman F., Bollars P., Donnelly M., Bellemans J., Nijs S.: Predictive factors for functional outcome and failure in angular stable osteosynthesis of the proximal humerus. *Injury.* 43: 153-158, 2012.
- 12) Cruess R.L.: Experience with steroid-induced avascular necrosis of the shoulder and etiologic considerations regarding osteonecrosis of the hip. *Clin Orthop.* 130: 86-93, 1978.
- 13) Beks RB., Ochen Y., Frima H., Smeeing DPJ., van der Meijden O., Timmers TK., van der Velde D., van Heijl M., Leenen LPH., Groenwold RHH., Houwert RM.: Operative versus nonoperative treatment of proximal humeral fractures: a systematic review, meta-analysis, and comparison of observational studies and randomized controlled trials. *J Shoulder Elbow Surg.* 27(8): 1526-1534, 2018.
- 14) Fjalestad T., Hole MØ., Hovden IA., Blücher J.,

- Strømsøe K.: Surgical treatment with an angular stable plate for complex displaced proximal humeral fractures in elderly patients: a randomized controlled trial. *J Orthop Trauma.* 26(2): 98-106, 2012.
- 15) Mauro E.C. Gracitelli., Eduardo A. Malavolta., Jorge H. Assuncao., Kodi E. Kojima., Paulo R. dos Reis., Jorge S. Silva., Arnaldo A. Ferreira Neto., Arnaldo J. Hernandez.: Locking intramedullary nails compared with locking plates for two- and three-part proximal humeral surgical neck fractures: a randomized controlled trial. *J Shoulder Elbow Surg.* 25(5): 695-703, 2016.
 - 16) Tamimi I., Montesa G., Collado F., González D., Carnero P., Rojas F., Nagib M., Pérez V., Álvarez M., Tamimi F.: Displaced proximal humeral fractures: when is surgery necessary? *Injury.* 46(10): 1921-1929, 2015.
 - 17) Boudard G., Pomares G., Milin L., Lemonnier I., Coudane H., Mainard D., Delagoutte JP.: Locking plate fixation versus antegrade nailing of 3- and 4-part proximal humerus fractures in patients without osteoporosis. Comparative retrospective study of 63 cases. *Orthop Traumatol Surg Res.* 100: 917-924, 2014.
 - 18) Konrad G., Audigé L., Lambert S., Hertel R., Südkamp NP.: Similar outcomes for nail versus plate fixation of three-part proximal humeral fractures. *Clin Orthop Relat Res.* 470(2): 602-609, 2011.
 - 19) Shintaro Y., Naoki S., Naomi O., Minami A.: Interlocking intramedullary nailing for nonunion of the proximal humerus with the Straight Nail System. *J Shoulder Elbow Surg.* 17: 755-759, 2008.
 - 20) Neer CS II.: Displaced proximal humeral fractures. II. Treatment of three-part and four-part displacement. *J Bone Joint Surg Am.* 52:1090-1103, 1970.
 - 21) Galatz LM., Iannotti JP.: Management of surgical neck nonunions. *Orthop Clin North Am.* 31: 51-61, 2000.
 - 22) Li M., Wang Y., Zhang Y., Yang M., Zhang P., Jiang B.: Intramedullary nail versus locking plate for treatment of proximal humeral fractures: A meta-analysis based on 1384 individuals. *J Int Med Res.* 46: 4363-4376, 2018.
 - 23) Hertel R., Hempfing A., Stiehler M., and Leunig M.: Predictors of humeral head ischemia after intracapsular fracture of the proximal humerus. *J Shoulder Elbow Surg.* 13: 427-433, 2004.
 - 24) Depalma AF.: *Surgery of the shoulder*, 3rd ed. JB Lippincott Co, Philadelphia. 348-427, 1983.
 - 25) Katsuya Nobuhara.: *Kata-sono kino to rinsho* second edition Igakushoin Tokyo. 275-286, 1987.
 - 26) Voigt C., Kreienborg S., Megatli O., Schulz AP., Lill H., Hurschler C.: How does a varus deformity of the humeral head affect elevation forces and shoulder function? A biomechanical study with human shoulder specimens. *J Orthop Trauma.* 25: 399-405, 2011.
 - 27) Führtmeier B., May R., Hente R., Maghsudi M., Völk M., Hammer J., Nerlich M., Prantl L.: Proximal humerus fractures: a comparative biomechanical analysis of intra and extramedullary implants. *Arch Orthop Trauma Surg.* 127(6): 441-447, 2007.
 - 28) Gardner MJ., Weil Y., Barker JU., Kelly BT., Helfet DL., Lorich DG.: The importance of medial support in locked plating of proximal humerus fractures. *J Orthop Trauma.* 21(3): 185-191, 2007.
 - 29) Park JY., An JW., Oh JH.: Open intramedullary nailing with tension band and locking sutures for proximal humeral fracture: Hot air balloon technique. *J Shoulder Elbow Surg.* 15(5): 594-601, 2006.
 - 30) Yoshioka C., Naoki S., Naomi O., Shintaro Y.: Complication of intramedullary nailing of proximal humerus fractures with the New Straight Nail System -Multicenter Study-. *Katakansetsu.* 36: 433-436, 2012.

